


JC17 Rec'd PCT/PTO 10 MAY 2001

| | | |
|---|--|---|
| FORM PTO-1390 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371 | | ATTORNEY'S DOCKET NUMBER PF-0643 USN |
| | | U.S. APPLICATION NO. (If known, see 37 CFR 1.5) TO BE ASSIGNED 097/831805 |
| INTERNATIONAL APPLICATION NO. PCT/US99/27566 | INTERNATIONAL FILING DATE 19 November 1999 | PRIORITY DATE CLAIMED 19 November 1998 |
| TITLE OF INVENTION IMMUNOGLOBULIN SUPERFAMILY PROTEINS | | |
| APPLICANT(S) FOR DO/EO/US INCYTE PHARMACEUTICALS, INC.; YUE, Henry; TANG, Y. Tom; CORLEY, Neil C.; GUEGLER, Karl J.; GORGONE, Gina A.; BAUGHN, Mariah R.; LU, Dyung Aina M.; LAL, Preeti; HILLMAN, Jennifer L.; YANG, Junming | | |
| Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: 1. <input checked="" type="checkbox"/> This is the FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input type="checkbox"/> This is an express request to promptly begin national examination procedures (35 U.S.C. 371 (f)). 4. <input type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (PCT Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau) b. <input type="checkbox"/> has been communicated by the International Bureau. c. <input checked="" type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). | | |
| Items 11 to 16 below concern document(s) or information included: | | |
| 11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.27 and 3.31 is included. 13. <input type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 14. <input type="checkbox"/> A substitute specification. 15. <input type="checkbox"/> A change of power of attorney and/or address letter. 16. <input checked="" type="checkbox"/> Other items or information: 1) Transmittal Letter (2 pp, in duplicate) 2) Return Postcard 3) Express Mail Label No.: <u>EL 856 113 141 US</u> 4) Request to Transfer | | |

RECEIVED

| | | | | | |
|---|--------------|--|------------|---|----|
| U.S. APPLICATION NO. (if known, see 37 CFR 1.5) TO BE ASSIGNED | | INTERNATIONAL APPLICATION NO.: PCT/US99/27566 | | ATTORNEY'S DOCKET NUMBER PF-0643 USN | |
| 17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO.....\$1000.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO..\$860.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO.....\$710.00 <input checked="" type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4).....\$690.00 <input type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4).....\$100.00 | | | | | |
| ENTER APPROPRIATE BASIC FEE AMOUNT = | | | | \$690.00 | |
| Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)). | | | | \$ | |
| CLAIMS | NUMBER FILED | NUMBER EXTRA | RATE | | |
| Total Claims | 20 = | 0 | X \$ 18.00 | \$ | |
| Independent Claims | 2 = | 0 | X \$ 80.00 | \$ | |
| MULTIPLE DEPENDENT CLAIM(S) (if applicable) | | | + \$270.00 | \$ | |
| TOTAL OF ABOVE CALCULATIONS = | | | | \$690.00 | |
| <input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2. | | | | \$ | |
| SUBTOTAL = | | | | \$690.00 | |
| Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)). | | | | \$ | |
| TOTAL NATIONAL FEE = | | | | \$690.00 | |
| Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by the appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property | | | | \$ | |
| TOTAL FEES ENCLOSED = | | | | \$690.00 | |
| | | | | Amount to be Refunded: | \$ |
| | | | | Charged: | \$ |
| a. <input type="checkbox"/> A check in the amount of \$_____ to cover the above fees is enclosed. b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. <u>09-0108</u> in the amount of \$ <u>690.00</u> to cover the above fees. c. <input type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>09-0108</u> . A duplicate copy of this sheet is enclosed. | | | | | |
| NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status. | | | | | |
| SEND ALL CORRESPONDENCE TO: INCYTE GENOMICS, INC. 3160 Porter Drive Palo Alto, CA 94304 | | | | | |
| SIGNATURE:  NAME: Diana Hamlet-Cox | | | | | |
| REGISTRATION NUMBER: 33,302 | | | | | |
| DATE: <u>10</u> May 2001 | | | | | |

T00T50: 508T2200

IMMUNOGLOBULIN SUPERFAMILY PROTEINS**TECHNICAL FIELD**

This invention relates to nucleic acid and amino acid sequences of immunoglobulin superfamily proteins and to the use of these sequences in the diagnosis, treatment, and prevention of cancer, immune system disorders, and infections.

5

BACKGROUND OF THE INVENTION

All vertebrates have developed sophisticated and complex immune systems that provide protection from viral, bacterial, fungal, and parasitic infections. Protection is mediated through cell surface and soluble molecules which function in recognition, adhesion or binding. The vertebrate immune system evolved from a common evolutionary precursor (i.e., these proteins have structural homology). A number of molecules outside the immune system that have similar functions are also derived from this same evolutionary precursor.

An important characteristic of the immune system is its ability to recognize and destroy foreign molecules, or antigens. Antigen recognition is mediated primarily by secreted and transmembrane proteins expressed by leukocytes (white blood cells) such as granulocytes, monocytes, and lymphocytes. Most of these proteins belong to the immunoglobulin (Ig) superfamily. The cell surface and soluble molecules of the immune system are classified as members of the Ig superfamily, members of which contain one or more repeats of a conserved structural Ig domain. The Ig domain, 70-110 amino acid residues in length, is homologous to either Ig variable-like (V) or Ig constant-like (C) domains. The Ig domain is described as antiparallel β sheets joined by a disulfide bond in an arrangement called the Ig fold. Members of the Ig superfamily include antibodies (Ab), T cell receptors (TCRs), class I and II major histocompatibility (MHC) proteins, CD2, CD3, CD4, CD8, poly-Ig receptors, Fc receptors, neural cell adhesion molecule (NCAM) and platelet-derived growth factor receptor (PDGFR).

Ig domains (V and C) are regions of conserved amino acid residues that give a polypeptide a globular tertiary structure called an immunoglobulin (or antibody) fold, which consists of two approximately parallel layers of β -sheets. Conserved cysteine residues form an intrachain disulfide-bonded loop, 55-75 amino acid residues in length, which connects the two layers of the β -sheets. Each β -sheet has three or four anti-parallel β -strands of 5-10 amino acid residues. Hydrophobic and hydrophilic interactions of amino acid residues within the β -strands stabilize the Ig fold (hydrophobic on inward facing amino acid residues and hydrophilic on the amino acid residues in the outward facing portion of the strands). A V domain consists of a longer polypeptide than a C domain, with an

additional pair of β -strands in the Ig fold.

A consistent feature of Ig superfamily genes is that each sequence of an Ig domain is encoded by a single exon. It is possible that the superfamily evolved from a gene coding for a single Ig domain involved in mediating cell-cell interactions. New members of the superfamily then arose by exon and gene duplications. Modern Ig superfamily proteins contain different numbers of V and/or C domains. Another evolutionary feature of this superfamily is the ability to undergo DNA rearrangements, a unique feature retained by the antigen receptor members of the family.

Many members of the Ig superfamily are integral plasma membrane proteins with extracellular Ig domains. The hydrophobic amino acid residues of their transmembrane domains and their cytoplasmic tails are very diverse, with little or no homology among Ig family members or to known signal-transducing structures. There are exceptions to this general superfamily description. For example, the cytoplasmic tail of PDGFR has tyrosine kinase activity. In addition Thy-1 is a glycoprotein found on thymocytes and T cells. This protein has no cytoplasmic tail, but is instead attached to the plasma membrane by a covalent glycosphosphatidylinositol linkage.

Another common feature of many Ig superfamily proteins is the interactions between Ig domains which are essential for the function of these molecules. Interactions between Ig domains of a multimeric protein can be either homophilic or heterophilic (i.e., between the same or different Ig domains). Antibodies are multimeric proteins which have both homophilic and heterophilic interactions between Ig domains. Pairing of constant regions of heavy chains forms the Fc region of an antibody and pairing of variable regions of light and heavy chains forms the antigen binding site of an antibody. Heterophilic interactions also occur between Ig domains of different molecules. These interactions provide adhesion between cells for significant cell-cell interactions in the immune system and in the developing and mature nervous system. (Reviewed in Abbas, A.K. et al. (1991) Cellular and Molecular Immunology, W.B. Saunders Company, Philadelphia, PA, pp.142-145.)

Antibodies

Antibodies, or immunoglobulins, are the founding members of the Ig superfamily and are the central components of the humoral immune response. Antibodies are either expressed on the surface of B-cells or secreted by B-cells into the circulation. Antibodies bind and neutralize blood-borne foreign antigens. The prototypical antibody is a tetramer consisting of two identical heavy polypeptide chains (H-chains) and two identical light polypeptide chains (L-chains) interlinked by disulfide bonds. This arrangement confers the characteristic Y-shape to antibody molecules. Antibodies are classified based on their H-chain composition. The five antibody classes, IgA, IgD, IgE, IgG and IgM, are defined by the α , δ , ϵ , γ , and μ H-chain types. There are two types of L-chains, κ and λ , either of which may associate as a pair with any H-chain pair. IgG, the most common class

of antibody found in the circulation, is tetrameric, while the other classes of antibodies are generally variants or multimers of this basic structure.

H-chains and L-chains each contain an N-terminal variable region and a C-terminal constant region. The constant region consists of about 110 amino acids in L-chains and about 330 or 440 amino acids in H-chains. The amino acid sequence of the constant region is nearly identical among H- or L-chains of a particular class. The variable region consists of about 110 amino acids in both H- and L-chains. However, the amino acid sequence of the variable region differs among H- or L-chains of a particular class. Within each H- or L-chain variable region are three hypervariable regions of extensive sequence diversity, each consisting of about 5 to 10 amino acids. In the antibody molecule, the H- and L-chain hypervariable regions come together to form the antigen recognition site. (Reviewed in Alberts, B. et al. (1994) Molecular Biology of the Cell, Garland Publishing, New York, NY, pages 1206-1213 and 1216-1217.)

Both H-chains and L-chains contain repeated Ig domains. For example, a typical H-chain contains four Ig domains, three of which occur within the constant region and one of which occurs within the variable region and contributes to the formation of the antigen recognition site. Likewise, a typical L-chain contains two Ig domains, one of which occurs within the constant region and one of which occurs within the variable region. In addition, H chains such as μ have been shown to associate with other polypeptides during differentiation of the B-cell. One such polypeptide called 8HS-20 is itself a member of the Ig superfamily and contains a single Ig domain (Shirasawa, T. et al. (1993) EMBO J. 12:1827-1834).

The immune system is capable of recognizing and responding to any foreign molecule that enters the body. Therefore, the immune system must be armed with a full repertoire of antibodies against all potential antigens. Such antibody diversity is generated by somatic rearrangement of gene segments encoding variable and constant regions. These gene segments are joined together by site-specific recombination which occurs between highly conserved DNA sequences that flank each gene segment. Because there are hundreds of different gene segments, millions of unique genes can be generated combinatorially. In addition, imprecise joining of these segments and an unusually high rate of somatic mutation within these segments further contribute to the generation of a diverse antibody population.

Antibodies can also be described in terms of their two main functional domains. Antigen recognition is mediated by the Fab (antigen binding fragment) region of the antibody, while effector functions are mediated by the Fc (crystallizable fragment) region. Binding of antibody to an antigen, such as a bacterium, triggers the destruction of the antigen by phagocytic white blood cells such as macrophages and neutrophils. These cells express surface receptors that specifically bind to the

antibody Fc region and allow the phagocytic cells to engulf, ingest, and degrade the antibody-bound antigen. The Fc receptors expressed by phagocytic cells are single-pass transmembrane glycoproteins of about 300 to 400 amino acids (Sears, D. W. et al. (1990) J. Immunol. 144:371-378). The extracellular portion of the Fc receptor typically contains two or three Ig domains.

- 5 Unique variants of Fc receptors have been identified in myeloid and lymphoid cells (Samaridis, J. and Colonna, M. (1997) Eur. J. Immunol. 27:660-665). Like typical Fc receptors, these proteins contain extracellular Ig domains and are encoded by cDNAs designated ILT1 and ILT2 (Ig-like transcripts 1 and 2). However, the transmembrane and cytoplasmic domains diverge significantly. In particular, the cytoplasmic domain is extended and contains protein motifs consistent
10 with a role in intracellular signal transduction.

- A new member of the Ig superfamily appears to play a structural role in the control of monocyte migration across epithelium or endothelium to sites of inflammation. This protein, called junctional adhesion molecule (JAM), is situated at tight junctions which occur between adjacent epithelial or endothelial cells (Martin-Padura, I. et al. (1998) J. Cell Biol. 142:117-127). JAM is 300
15 amino acids in length and contains two Ig domains. A monoclonal antibody (mAb) directed against JAM inhibited transmigration of monocytes across endothelial cell layers in vitro. Furthermore, systemic administration of this mAb to mice prevented recruitment of monocytes to sites of inflammation.

- Viral proteins which contain Ig domains have also been described (Senkevich, T. G. et al. (1996) Science 273:813-816). These proteins include MHC-like homologs identified in the human tumorigenic poxvirus Molluscum contagiosum. Such proteins may provide a mechanism by which
20 the virus evades the host's immunologic surveillance system.

- T-cell receptors are both structurally and functionally related to antibodies. (Reviewed in Alberts, supra, pp. 1228-1229.) T-cell receptors are cell surface proteins that bind foreign antigens
25 and mediate diverse aspects of the immune response. A typical T-cell receptor is a heterodimer comprised of two disulfide-linked polypeptide chains called α and β . Each chain is about 280 amino acids in length and contains one variable region and one constant region. Each variable or constant region folds into an Ig domain. The variable regions from the α and β chains come together in the heterodimer to form the antigen recognition site. T-cell receptor diversity is generated by somatic
30 rearrangement of gene segments encoding the α and β chains. T-cell receptors recognize small peptide antigens that are expressed on the surface of antigen-presenting cells and pathogen-infected cells. These peptide antigens are presented on the cell surface in association with major histocompatibility proteins which provide the proper context for antigen recognition.

Synaptic Membrane Glycoproteins

Specialized cell junctions can occur at points of cell-cell contact. Among these cell junctions are communicating junctions which mediate the passage of chemical and electrical signals between cells. In the central nervous system, communicating junctions between neurons are known as synaptic junctions. They are composed of the membranes and cytoskeletons of the pre- and post-synaptic neurons. Some glycoproteins, found in biochemically isolated synaptic subfractions such as the synaptic membrane (SM) and postsynaptic density (PSD) fractions, have been identified and their functions established. An example is the SM glycoprotein, gp50, identified as the $\beta 2$ subunit of the Na^+/K^+ -ATPase.

Glycoproteins in the SM and PSD which have their oligosaccharide domains facing the synaptic junction are in a good position to mediate adhesive interactions between neurons. The PAC 1 glycoproteins, components of the PSD, have been identified as members of the cadherin family, proteins involved in the Ca^{2+} -dependent cell-cell adhesion in vertebrate tissues. Further support of these molecules mediating adhesive interactions is the presence of integrin-type adhesion molecules and NCAM, a member of the Ig superfamily, in the SM.

Two glycoproteins, gp65 and gp55, are major components of synaptic membranes prepared from rat forebrain. They are members of the Ig superfamily containing three and two Ig domains, respectively. As members of the Ig superfamily, it is proposed that a possible function of these proteins is to mediate adhesive interactions at the synaptic junction. (Langnaese, K. et al. (1997) J. Biol. Chem. 272(2):821-827.)

The discovery of new immunoglobulin superfamily proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cancer, immune system disorders, and infections.

25 SUMMARY OF THE INVENTION

The invention features substantially purified polypeptides, immunoglobulin superfamily proteins, referred to collectively as "IGFAM" and individually as "IGFAM-1," "IGFAM-2," "IGFAM-3," "IGFAM-4," "IGFAM-5," "IGFAM-6," "IGFAM-7," "IGFAM-8," "IGFAM-9," "IGFAM-10," "IGFAM-11," "IGFAM-12," "IGFAM-13," "IGFAM-14," "IGFAM-15," "IGFAM-16," "IGFAM-17," "IGFAM-18," and "IGFAM-19." In one aspect, the invention provides a substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19 and fragments thereof. The invention also includes a polypeptide comprising an amino acid sequence that differs by one or more conservative amino acid substitutions from an amino acid sequence selected from the group consisting of SEQ ID NO:1-19.

The invention further provides a substantially purified variant having at least 90% amino acid identity to at least one of the amino acid sequences selected from the group consisting of SEQ ID NO:1-19 and fragments thereof. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of
5 SEQ ID NO:1-19 and fragments thereof. The invention also includes an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19 and fragments thereof.

Additionally, the invention provides an isolated and purified polynucleotide which hybridizes
10 under stringent conditions to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19 and fragments thereof. The invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1-19 and fragments thereof.

15 The invention also provides a method for detecting a polynucleotide in a sample containing nucleic acids, the method comprising the steps of: (a) hybridizing the complement of the polynucleotide sequence to at least one of the polynucleotides of the sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of a polynucleotide in the sample. In one aspect,
20 the method further comprises amplifying the polynucleotide prior to hybridization.

The invention also provides an isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38 and fragments thereof. The invention further provides an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide sequence selected from the group
25 consisting of SEQ ID NO:20-38 and fragments thereof. The invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38 and fragments thereof.

The invention further provides an expression vector containing at least a fragment of the
30 polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19. In another aspect, the expression vector is contained within a host cell.

The invention also provides a method for producing a polypeptide, the method comprising the steps of: (a) culturing the host cell containing an expression vector containing a polynucleotide of the

invention under conditions suitable for the expression of the polypeptide; and (b) recovering the polypeptide from the host cell culture.

The invention also provides a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-19 and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

The invention further includes a purified antibody which binds to a polypeptide selected from the group consisting of SEQ ID NO:1-19 and fragments thereof. The invention also provides a purified agonist and a purified antagonist to the polypeptide.

The invention also provides a method for treating or preventing a disorder associated with decreased expression or activity of IGFAM, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-19 and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

The invention also provides a method for treating or preventing a disorder associated with increased expression or activity of IGFAM, the method comprising administering to a subject in need of such treatment an effective amount of an antagonist of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-19 and fragments thereof.

BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding IGFAM.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of IGFAM.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding IGFAM were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze IGFAM, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing
5 particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a
10 reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be
15 used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

20 DEFINITIONS

"IGFAM" refers to the amino acid sequences of substantially purified IGFAM obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of
25 IGFAM. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of IGFAM either by directly interacting with IGFAM or by acting on components of the biological pathway in which IGFAM participates.

An "allelic variant" is an alternative form of the gene encoding IGFAM. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in
30 polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding IGFAM include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as IGFAM or a polypeptide with at least one functional characteristic of IGFAM. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding IGFAM, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding IGFAM. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent IGFAM. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of IGFAM is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where “amino acid sequence” is recited to refer to an amino acid sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

“Amplification” relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term “antagonist” refers to a molecule which inhibits or attenuates the biological activity of IGFAM. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of IGFAM either by directly interacting with IGFAM or by acting on components of the biological pathway in which IGFAM participates.

The term “antibody” refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind IGFAM polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or

oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize
 5 the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures
 10 on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition containing a nucleic acid sequence which is complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the
 15 complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the
 20 capability of the natural, recombinant, or synthetic IGFAM, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules
 25 may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid
 30 (PNA) molecules.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding IGFAM or fragments of IGFAM may

be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

5 "Consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using the XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of one or more Incyte Clones and, in some cases, one or more public domain ESTs and/or cDNAs, using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG,
10 Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that, when made, least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows
15 amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

| | Original Residue | Conservative Substitution |
|----|------------------|---------------------------|
| | Ala | Gly, Ser |
| | Arg | His, Lys |
| 20 | Asn | Asp, Gln, His |
| | Asp | Asn, Glu |
| | Cys | Ala, Ser |
| | Gln | Asn, Glu, His |
| | Glu | Asp, Gln, His |
| 25 | Gly | Ala |
| | His | Asn, Arg, Gln, Glu |
| | Ile | Leu, Val |
| | Leu | Ile, Val |
| | Lys | Arg, Gln, Glu |
| 30 | Met | Leu, Ile |
| | Phe | His, Met, Leu, Trp, Tyr |
| | Ser | Cys, Thr |
| | Thr | Ser, Val |
| | Trp | Phe, Tyr |
| 35 | Tyr | His, Phe, Trp |
| | Val | Ile, Leu, Thr |

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation,
40 (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

10 A "fragment" is a unique portion of IGFAM or the polynucleotide encoding IGFAM which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

20 A fragment of SEQ ID NO:20-38 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:20-38, for example, as distinct from any other sequence in the same genome. A fragment of SEQ ID NO:20-38 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:20-38 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:20-38 and the region of SEQ ID NO:20-38 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

25 A fragment of SEQ ID NO:1-19 is encoded by a fragment of SEQ ID NO:20-38. A fragment of SEQ ID NO:1-19 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-19. For example, a fragment of SEQ ID NO:1-19 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-19. The precise length of a fragment of SEQ ID NO:1-19 and the region of SEQ ID NO:1-19 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequence pairs.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis

programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The

5 "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

10 *Reward for match: 1*

Penalty for mismatch: -2

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

15 *Word Size: 11*

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at

20 least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode

25 similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a

30 standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the hydrophobicity and acidity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default

parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default
 5 residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) with blastp set at default parameters. Such default parameters may be, for example:

10 *Matrix: BLOSUM62*
 Open Gap: 11 and Extension Gap: 1 penalties
 Gap x drop-off: 50
 Expect: 10
 Word Size: 3
 15 *Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least
 20 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for
 25 stable mitotic chromosome segregation and maintenance.

The term "humanized antibody" refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a
 30 complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of identity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding

between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive

5 annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Generally, such wash temperatures are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic

10 strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

15 High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance,

20 denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative

25 of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid

30 support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune

disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

5 The terms "element" and "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term "modulate" refers to a change in the activity of IGFAM. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of IGFAM.

10 The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a
15 functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which
20 comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Probe" refers to nucleic acid sequences encoding IGFAM, their complements, or fragments
25 thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

"Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target
30 polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100,

or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis et al., 1990, PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence.

This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a
5 recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is
10 expressed, inducing a protective immunological response in the mammal.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding IGFAM, or fragments thereof, or IGFAM itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a
15 protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide
20 containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which
25 they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers,
30 microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

"Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic

acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an

5 autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-

10 1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of

15 polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene

20 between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having

25 at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

30

THE INVENTION

The invention is based on the discovery of new human immunoglobulin superfamily proteins (IGFAM), the polynucleotides encoding IGFAM, and the use of these compositions for the diagnosis, treatment, or prevention of cancer, immune system disorders, and infections.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding IGFAM. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each IGFAM were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. The Incyte clones in column 5 were used to assemble the consensus nucleotide sequence of each IGFAM and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs. Note that in column 5, all but one of the polypeptides of the invention contain one or more Ig domains as predicted by protein function analysis programs such as PROFILESCAN, BLIMPS, PFAM, and MOTIFS. The polypeptide of the invention which lacks a predicted Ig domain (SEQ ID NO:4) does, however, show significant similarity with the viral Ig-containing protein MC51L-53L-54L.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding IGFAM. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:20-38 and to distinguish between SEQ ID NO:20-38 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express IGFAM as a fraction of total tissues expressing IGFAM. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing IGFAM as a fraction of total tissues expressing IGFAM. Column 5 lists the vectors used to subclone each cDNA library. Note that the nucleotide sequences of SEQ ID NO:23, and SEQ ID NO:27 are expressed primarily in cells and tissues associated with the hematopoietic/immune system and inflammation.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding IGFAM were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3

shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

The invention also encompasses IGFAM variants. A preferred IGFAM variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the IGFAM amino acid sequence, and which contains at least one functional or structural characteristic of IGFAM.

The invention also encompasses polynucleotides which encode IGFAM. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:20-38, which encodes IGFAM.

The invention also encompasses a variant of a polynucleotide sequence encoding IGFAM. In particular, such a variant polynucleotide sequence will have at least about 80%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding IGFAM. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:20-38 which has at least about 80%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:20-38. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of IGFAM.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding IGFAM, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring IGFAM, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode IGFAM and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring IGFAM under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding IGFAM or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding IGFAM and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a

greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode IGFAM and IGFAM derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems
5 using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding IGFAM or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:20-38 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and
10 S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment
15 of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV),
20 PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Perkin-Elmer). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Perkin-Elmer), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short
25 Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding IGFAM may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed,
30 restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids*

Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown
5 sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using
10 commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been
15 size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze
20 the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process
25 from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode IGFAM may be cloned in recombinant DNA molecules that direct expression of
30 IGFAM, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express IGFAM.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter IGFAM-encoding sequences for a variety of purposes including, but

not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction
5 sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In another embodiment, sequences encoding IGFAM may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-232.) Alternatively, IGFAM itself or a fragment thereof may be synthesized using chemical methods. For
10 example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of IGFAM, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

15 The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY.)

20 In order to express a biologically active IGFAM, the nucleotide sequences encoding IGFAM or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in
25 polynucleotide sequences encoding IGFAM. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding IGFAM. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding IGFAM and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional
30 transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used.

(See, e.g., Scharf, D. et al. (1994) Results Probl. Cell Differ. 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding IGFAM and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding IGFAM. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding IGFAM. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding IGFAM can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding IGFAM into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of IGFAM are needed, e.g. for the production of antibodies, vectors which direct high level expression of IGFAM may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of IGFAM. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of IGFAM. Transcription of sequences encoding IGFAM may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock
5 promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

10 In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding IGFAM may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses IGFAM in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc.
15 Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are
20 constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of IGFAM in cell lines is preferred. For example, sequences encoding IGFAM can be transformed into
25 cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express
30 the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *ap^r* cells, respectively. (See, e.g., Wigler, M. et

al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g.,

5 Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate

10 luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the

15 sequence encoding IGFAM is inserted within a marker gene sequence, transformed cells containing sequences encoding IGFAM can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding IGFAM under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

20 In general, host cells that contain the nucleic acid sequence encoding IGFAM and that express IGFAM may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

25 Immunological methods for detecting and measuring the expression of IGFAM using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on IGFAM is preferred, but a

30 competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding IGFAM include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding IGFAM, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding IGFAM may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode IGFAM may be designed to contain signal sequences which direct secretion of IGFAM through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding IGFAM may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric IGFAM protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of IGFAM activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST),

maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity

- 5 purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the IGFAM encoding sequence and the heterologous protein sequence, so that IGFAM may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10).
- 10 A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled IGFAM may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the

- 15 T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

Fragments of IGFAM may be produced not only by recombinant means, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be

- 20 achieved, for example, using the ABI 431A peptide synthesizer (Perkin-Elmer). Various fragments of IGFAM may be synthesized separately and then combined to produce the full length molecule.

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of IGFAM and immunoglobulin superfamily proteins and Ig domain-containing proteins such

- 25 as antibody heavy and light chains. In addition, the expression of IGFAM is closely associated with proliferating tissues, cancerous tissue and with hematopoiesis, inflammation, and other processes mediated by the immune system. Therefore, IGFAM appears to play a role in cancer, immune system disorders, and infections. In the treatment of disorders associated with increased IGFAM expression or activity, it is desirable to decrease the expression or activity of IGFAM. In the treatment of
- 30 disorders associated with decreased IGFAM expression or activity, it is desirable to increase the expression or activity of IGFAM.

Therefore, in one embodiment, IGFAM or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of IGFAM. Examples of such disorders include, but are not limited to, a cancer such as

- adenocarcinoma, melanoma, sarcoma, teratocarcinoma, and in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a disorder of the immune system such as inflammation,
- 5 actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythroblastosis fetalis, erythema nodosum, atrophic gastritis,
- 10 glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome, episodic lymphopenia with lymphocytotoxins, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid
- 15 arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; and an infection caused by a viral agent classified as adenovirus, arenavirus, bunyavirus, calicivirus, coronavirus, filovirus,
- 20 hepadnavirus, herpesvirus, flavivirus, orthomyxovirus, parvovirus, papovavirus, paramyxovirus, picornavirus, poxvirus, reovirus, retrovirus, rhabdovirus, or togavirus; an infection caused by a bacterial agent classified as pneumococcus, staphylococcus, streptococcus, bacillus, corynebacterium, clostridium, meningococcus, gonococcus, listeria, moraxella, kingella, haemophilus, legionella, bordetella, gram-negative enterobacterium including shigella, salmonella, or campylobacter,
- 25 pseudomonas, vibrio, brucella, francisella, yersinia, bartonella, norcardium, actinomyces, mycobacterium, spirochaetale, rickettsia, chlamydia, or mycoplasma; an infection caused by a fungal agent classified as aspergillus, blastomyces, dermatophytes, cryptococcus, coccidioides, malassezia, histoplasma, or other mycosis-causing fungal agent; and an infection caused by a parasite classified as plasmodium or malaria-causing, parasitic entamoeba, leishmania, trypanosoma, toxoplasma,
- 30 pneumocystis carinii, intestinal protozoa such as giardia, trichomonas, tissue nematode such as trichinella, intestinal nematode such as ascaris, lymphatic filarial nematode, trematode such as schistosoma, and cestode such as tapeworm.

In another embodiment, a vector capable of expressing IGFAM or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased

expression or activity of IGFAM including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified IGFAM in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of IGFAM including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of IGFAM may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of IGFAM including, but not limited to, those listed above.

In a further embodiment, an antagonist of IGFAM may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of IGFAM. Examples of such disorders include, but are not limited to, those cancer, immune system disorders, and infections described above. In one aspect, an antibody which specifically binds IGFAM may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express IGFAM.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding IGFAM may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of IGFAM including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of IGFAM may be produced using methods which are generally known in the art. In particular, purified IGFAM may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind IGFAM. Antibodies to IGFAM may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with IGFAM or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to

increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

- 5 It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to IGFAM have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of IGFAM
- 10 amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

- Monoclonal antibodies to IGFAM may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma
- 15 technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

- In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate
- 20 antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce IGFAM-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be
- 25 generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

- Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA
- 30 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for IGFAM may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and

easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between IGFAM and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering IGFAM epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for IGFAM. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of IGFAM-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple IGFAM epitopes, represents the average affinity, or avidity, of the antibodies for IGFAM. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular IGFAM epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the IGFAM-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of IGFAM, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington, DC; Liddell, J.E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of IGFAM-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding IGFAM, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding IGFAM may be used in situations in which it would be

desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding IGFAM. Thus, complementary molecules or fragments may be used to modulate IGFAM activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments
5 can be designed from various locations along the coding or control regions of sequences encoding IGFAM.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used
10 to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding IGFAM. (See, e.g., Sambrook, supra; Ausubel, 1995, supra.)

Genes encoding IGFAM can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding IGFAM. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in
15 the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing
20 complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding IGFAM. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may be employed. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the
25 binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

30 Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding IGFAM.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for
5 secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques
10 for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding IGFAM. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into
15 cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be
20 extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use *in vivo*, *in vitro*, and *ex vivo*. For *ex vivo* therapy, vectors may be introduced into stem cells
25 taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of
30 such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of IGFAM,

antibodies to IGFAM, and mimetics, agonists, antagonists, or inhibitors of IGFAM. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol.

Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

5 Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily
10 injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

15 For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

 The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

20 The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a
25 pH range of 4.5 to 5.5, that is combined with buffer prior to use.

 After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of IGFAM, such labeling would include amount, frequency, and method of administration.

 Pharmaceutical compositions suitable for use in the invention include compositions wherein
30 the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

 For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of

administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example IGFAM or fragments thereof, antibodies of IGFAM, and agonists, antagonists or inhibitors of IGFAM, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind IGFAM may be used for the diagnosis of disorders characterized by expression of IGFAM, or in assays to monitor patients being treated with IGFAM or agonists, antagonists, or inhibitors of IGFAM. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for IGFAM include methods which utilize the antibody and a label to detect

IGFAM in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

5 A variety of protocols for measuring IGFAM, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of IGFAM expression. Normal or standard values for IGFAM expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to IGFAM under conditions suitable for complex formation. The amount of standard complex formation
10 may be quantitated by various methods, such as photometric means. Quantities of IGFAM expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding IGFAM may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences,
15 complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of IGFAM may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of IGFAM, and to monitor regulation of IGFAM levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide
20 sequences, including genomic sequences, encoding IGFAM or closely related molecules may be used to identify nucleic acid sequences which encode IGFAM. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding IGFAM, allelic variants, or related
25 sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the IGFAM encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:20-38 or from genomic sequences including promoters, enhancers, and introns of the IGFAM gene.

30 Means for producing specific hybridization probes for DNAs encoding IGFAM include the cloning of polynucleotide sequences encoding IGFAM or IGFAM derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a

variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding IGFAM may be used for the diagnosis of disorders associated with expression of IGFAM. Examples of such disorders include, but are not limited to, a

5 cancer such as adenocarcinoma, melanoma, sarcoma, teratocarcinoma, and in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a disorder of the immune system such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease,

10 adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's

15 thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome, episodic lymphopenia with lymphocytotoxins, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic

20 lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; and an infection caused by a viral agent classified as adenovirus, arenavirus, bunyavirus, calicivirus, coronavirus, filovirus, hepadnavirus, herpesvirus, flavivirus, orthomyxovirus, parvovirus,

25 papovavirus, paramyxovirus, picornavirus, poxvirus, reovirus, retrovirus, rhabdovirus, or togavirus; an infection caused by a bacterial agent classified as pneumococcus, staphylococcus, streptococcus, bacillus, corynebacterium, clostridium, meningococcus, gonococcus, listeria, moraxella, kingella, haemophilus, legionella, bordetella, gram-negative enterobacterium including shigella, salmonella, or campylobacter, pseudomonas, vibrio, brucella, francisella, yersinia, bartonella, norcardium,

30 actinomyces, mycobacterium, spirochaetale, rickettsia, chlamydia, or mycoplasma; an infection caused by a fungal agent classified as aspergillus, blastomyces, dermatophytes, cryptococcus, coccidioides, malassezia, histoplasma, or other mycosis-causing fungal agent; and an infection caused by a parasite classified as plasmodium or malaria-causing, parasitic entamoeba, leishmania, trypanosoma, toxoplasma, pneumocystis carinii, intestinal protozoa such as giardia, trichomonas,

tissue nematode such as trichinella, intestinal nematode such as ascaris, lymphatic filarial nematode, trematode such as schistosoma, and cestode such as tapeworm. The polynucleotide sequences encoding IGFAM may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in
5 microarrays utilizing fluids or tissues from patients to detect altered IGFAM expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding IGFAM may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding IGFAM may be labeled by standard methods and added to a fluid or tissue
10 sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding IGFAM in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate
15 the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of IGFAM, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a
20 sequence, or a fragment thereof, encoding IGFAM, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard
25 values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several
30 days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals

to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding IGFAM may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced *in vitro*. Oligomers will preferably contain a fragment of a polynucleotide encoding IGFAM, or a fragment of a polynucleotide complementary to the polynucleotide encoding IGFAM, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

Methods which may also be used to quantify the expression of IGFAM include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

In another embodiment of the invention, nucleic acid sequences encoding IGFAM may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355; Price, C.M. (1993) *Blood Rev.* 7:127-134; and Trask, B.J. (1991) *Trends Genet.* 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the
5 location of the gene encoding IGFAM on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as
10 linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other
15 gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among
20 normal, carrier, or affected individuals.

In another embodiment of the invention, IGFAM, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes
25 between IGFAM and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with IGFAM, or fragments thereof,
30 and washed. Bound IGFAM is then detected by methods well known in the art. Purified IGFAM can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing

antibodies capable of binding IGFAM specifically compete with a test compound for binding IGFAM. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with IGFAM.

In additional embodiments, the nucleotide sequences which encode IGFAM may be used in
5 any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are,
10 therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific
15 embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. [Attorney Docket No. PF-0643 P, filed November 19, 1998], U.S. Ser. No. 60/113,635, and U.S. Ser. No. 60/128,194, are hereby expressly incorporated by reference.

20

EXAMPLES

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a
25 monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A⁺) RNA was isolated
30 using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA

libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic
5 oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g.,
10 PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), or pINCY (Incyte Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

15 Plasmids were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN.
20 Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in
25 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

cDNA sequencing reactions were processed using standard methods or high-throughput
30 instrumentation such as the ABI CATALYST 800 (Perkin-Elmer) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Perkin-Elmer) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading
5 frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example V.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the
10 art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other
15 parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence
20 alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried
25 against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA.
30 The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene

families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:20-38. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

100

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding IGFAM occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Extension of IGFAM Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:20-38 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this

fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target
5 sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR
10 was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C,
15 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN
20 quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by
25 electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For
30 shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on

antibiotic-containing media. individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

In like manner, the nucleotide sequences of SEQ ID NO:20-38 are used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

VI. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:20-38 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

VII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, *supra*.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

VIII. Complementary Polynucleotides

Sequences complementary to the IGFAM-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring IGFAM. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of IGFAM. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the IGFAM-encoding transcript.

IX. Expression of IGFAM

Expression and purification of IGFAM is achieved using bacterial or virus-based expression systems. For expression of IGFAM in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid

promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3).

Antibiotic resistant bacteria express IGFAM upon induction with isopropyl beta-D-

thiogalactopyranoside (IPTG). Expression of IGFAM in eukaryotic cells is achieved by infecting

5 insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding IGFAM by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to
10 infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, IGFAM is synthesized as a fusion protein with, e.g., glutathione
15 S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from
20 IGFAM at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified IGFAM obtained by these methods can be used directly in the
25 following activity assay.

X. Demonstration of IGFAM Activity

An assay for IGFAM activity measures the ability of IGFAM to recognize and precipitate antigens from serum. This activity can be measured by the quantitative precipitin reaction. (Golub, E. S. et al. (1987) Immunology: A Synthesis, Sinauer Associates, Sunderland, MA, pages 113-115.)

30 IGFAM is isotopically labeled using methods known in the art. Various serum concentrations are added to constant amounts of labeled IGFAM. IGFAM-antigen complexes precipitate out of solution and are collected by centrifugation. The amount of precipitable IGFAM-antigen complex is proportional to the amount of radioisotope detected in the precipitate. The amount of precipitable IGFAM-antigen complex is plotted against the serum concentration. For various serum

concentrations, a characteristic precipitin curve is obtained, in which the amount of precipitable IGFAM-antigen complex initially increases proportionately with increasing serum concentration, peaks at the equivalence point, and then decreases proportionately with further increases in serum concentration. Thus, the amount of precipitable IGFAM-antigen complex is a measure of IGFAM activity which is characterized by sensitivity to both limiting and excess quantities of antigen.

Alternatively, an assay for IGFAM activity measures the expression of IGFAM on the cell surface. cDNA encoding IGFAM is transfected into a non-leukocytic cell line. Cell surface proteins are labeled with biotin (de la Fuente, M.A. et.al. (1997) Blood 90:2398-2405). Immunoprecipitations are performed using IGFAM-specific antibodies, and immunoprecipitated samples are analyzed using SDS-PAGE and immunoblotting techniques. The ratio of labeled immunoprecipitant to unlabeled immunoprecipitant is proportional to the amount of IGFAM expressed on the cell surface.

XI. Functional Assays

IGFAM function is assessed by expressing the sequences encoding IGFAM at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT (Life Technologies) and pCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of IGFAM on gene expression can be assessed using highly purified

populations of cells transfected with sequences encoding IGFAM and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding IGFAM and other genes of interest can be analyzed by northern analysis or microarray techniques.

XII. Production of IGFAM Specific Antibodies

IGFAM substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the IGFAM amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-IGFAM activity by, for example, binding the peptide or IGFAM to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIII. Purification of Naturally Occurring IGFAM Using Specific Antibodies

Naturally occurring or recombinant IGFAM is substantially purified by immunoaffinity chromatography using antibodies specific for IGFAM. An immunoaffinity column is constructed by covalently coupling anti-IGFAM antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing IGFAM are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of IGFAM (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/IGFAM binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and IGFAM is collected.

XIV. Identification of Molecules Which Interact with IGFAM

IGFAM, or biologically active fragments thereof, are labeled with ^{125}I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) Biochem. J. 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled IGFAM, washed, and any wells with labeled IGFAM complex are assayed. Data obtained using different concentrations of IGFAM are used to calculate values for the number, affinity, and association of IGFAM with the candidate molecules.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

| Polypeptide SEQ ID NO: | Nucleotide SEQ ID NO: | Clone ID | Library | Fragments | |
|---------------------------|--------------------------|----------|-----------|--|--|
| | | | | | |
| 1 | 20 | 079785 | SYNORAB01 | 079785H1 (SYNORAB01), 079785R6 (SYNORAB01), 930708R1 (CERVNOT01) | |
| 2 | 21 | 2469025 | THYRN0T08 | 731672R1 (LUNGNOT03), 1241709R6 (LUNGNOT03), 1535284H1 (SPLNNOT04), 2242944T6 (PANCUT02), 2354183H1 (LUNGNOT20), 2469025H1 (THYRN0T08), 3370185H1 (CONNTUT04), 3508954H1 (CONCNOT01), 3882135F6 (SPLNNOT11), 5021181H1 (OVARNON03) | |
| 3 | 22 | 2906265 | THYMN0T05 | 2684470T6 (LUNGNOT23), 2906265H1 (THYMN0T05), 2906265T6 (THYMN0T05), SBQA01965D1 | |
| 4 | 23 | 788975 | PROSTUT03 | 334315R6 (EOSIHET02), 788975H1 (PROSTUT03), 2995078H1 (OVARTUT07), 3210491T6 (BLADNOT08), SAJA03793F1, SAJA00460R1 | |
| 5 | 24 | 1407148 | LATRTUT02 | 1407148H1 (LATRTUT02), 1666491T6 (BMARNOT03), 3339721F6 (SPLNNOT10) | |
| 6 | 25 | 1870848 | SKINBIT01 | 1600994F6 (BLADNOT03), 1870848F6 (SKINBIT01), 1870848H1 (SKINBIT01), 1870848T3 (SKINBIT01), 3586454H1 (293TF4T01), SBGA04741F1, SBGA05485F1 | |
| 7 | 26 | 1888468 | BLADTUT07 | 1888468H1 (BLADTUT07), 2848829H1 (BRSTTUT13), 3336345F6 (SPLNNOT10) | |
| 8 | 27 | 2770104 | COLANOT02 | 034758X22 (THP1NOB01), 034758X34R1 (THP1NOB01), 169537X14 (BMARNOR02), 1526466F1 (UCMCL5T01), 2770104H1 (COLANOT02) | |
| 9 | 28 | 2851053 | BRSTTUT13 | 540284R1 (LNODNOT02), 689572R6 (LUNGUT02), 2851053H1 (BRSTTUT13) | |

Table 1 cont.

| Polypeptide SEQ ID NO: | Nucleotide SEQ ID NO: | Clone ID | Library | Fragments |
|---------------------------|--------------------------|----------|-----------|--|
| 10 | 29 | 3238787 | COLAUCT01 | 540284F1 (LNODNOT02), 792219T1 (PROSTUT03), 2179956H1 (SININOT01), 2630239T6 (COLNTUT15), 3238787H1 (COLAUCT01) |
| 11 | 30 | 3559548 | LUNGNOT31 | 701560H1 (SYNORAT03), 3336345F6 (SPLNNOT10), 3559548H1 (LUNGNOT31) |
| 12 | 31 | 3872741 | BMARNOT03 | 2354696F6 (LUNGNOT20), 2998376F6 (OVARTUT07), 3872741H1 (BMARNOT03) |
| 13 | 32 | 3981428 | LUNGTUT08 | 930366T1 (CERVNOT01), 3981428F6 (LUNGTUT08), 3981428H1 (LUNGTUT08) |
| 14 | 33 | 4635039 | GBLADIT02 | 2849560F6 (BRSTTUT13), 4635039H1 (GBLADIT02) |
| 15 | 34 | 3240710 | COLAUCT01 | 1820516T6 (GBLATUT01), 3240710H1 (COLAUCT01), 3344660F6 (SPLNNOT09) |
| 16 | 35 | 4945813 | SINTNOT25 | 1435848F6 (PANCNOT08), 1822478F6 (GBLATUT01), 4945813H1 (SINTNOT25), SBJA01188F1, SBPA02549D1 |
| 17 | 36 | 4948957 | SINTNOT25 | 2183186H1 (SININOT01), 4948957H1 (SINTNOT25), SAWA00061F1 |
| 18 | 37 | 4949649 | SINTNOT25 | 232824F1 (SINTNOT02), 991833T1 (COLNNOT11), 2994953H1 (OVARTUT07), 3860911T6 (LNODNOT03), 4949649H1 (SINTNOT25) |
| 19 | 39 | 5500302 | BRABDIR01 | 036351X10, 036351X6 and 036351X9 (HUVENOB01), 777804R1 (COLNNOT05), 810532R1 (LUNGNOT04), 2538351H1 (BONRTUT01), 5500302H1 (BRABDIR01) |

Table 2

| Polypeptide SEQ ID NO: | Amino Acid Residues | Potential Phosphorylation Sites | Potential Glycosylation Sites | Signature Sequences Motifs, and Domains | Homologous Sequences | Analytical Methods and Databases |
|---------------------------|---------------------------|---|-------------------------------------|--|--|---|
| 1 | 237 | S36 S89 T125 S185 T187 S205 T44 | | Ig domains G38-Q112 S150-V219 D193-E236 Ig signatures S154-A176 Y215-F232 Signal peptide M1-C22 | κ L-chain (g2765423) (Homo sapiens) | BLAST MOTIFS PFAM PROFILES SCAN BLOCKS HMM SPSCAN |
| 2 | 537 | S215 T117 T206 S246 S511 S93 T104 S110 S256 T284 T288 Y529 | N83 N378 N469 N520 | Ig domains S41-R128 N174-V239 R336-L404 E440-V510 T486-M536 Ig signatures Y235-H241 W444-R466 Y506-R523 Signal peptide M1-S26 | δ H-chain (g495871) (Homo sapiens) | BLAST MOTIFS PFAM PROFILES SCAN BLOCKS SPSCAN |

Table 2 cont.

| Polypeptide SEQ ID NO: | Amino Acid Residues | Potential Phosphorylation Sites | Potential Glycosylation Sites | Signature Sequences Motifs, and Domains | Homologous Sequences | Analytical Methods and Databases |
|---------------------------|---------------------------|--|-------------------------------------|--|---|--|
| 3 | 311 | S72 T101 T132 S151 S238 S28 T32 S81 T158 S211 | N204 | Ig domains G35-S113 T158-S211 Ig signatures T162-K184 F228-Q245 Signal peptide M1-A21 Transmembrane domain Y283-M303 | T cell receptor β chain (g3002935) (Homo sapiens) | BLAST MOTIFS PFAM BLOCKS HMM SPSCAN |
| 4 | 194 | S46 T70 S105 T119 T2 S43 | N79 N94 N103 N147 | Signal peptide M1-A30 | MC51L-53L-54L homolog (secreted glycoprotein) (g5231020) (Molluscum contagiosum virus) | BLAST MOTIFS |
| 5 | 236 | S36 T124 S184 T186 S204 T44 | | Ig domains G38-Q112 S149-V218 D192-E235 Signal peptide M1-C22 | Ig κ light chain (g2765423) (Homo sapiens) | BLAST MOTIFS PROFILES BLOCKS PFAM SPSCAN HMM |

Table 2 cont.

| Polypeptide SEQ ID NO: | Amino Acid Residues | Potential Phosphorylation Sites | Potential Glycosylation Sites | Signature Sequences Motifs, and Domains | Homologous Sequences | Analytical Methods and Databases |
|---------------------------|---------------------------|--|-------------------------------------|--|--|--|
| 6 | 310 | S110 T106 T296 S37 S99 S227 S281 Y77 | N104 N192 | Ig domains F46-V117 G153-A221 Signal peptide M1-G30 Transmembrane domain: Y238-T260 | JAM: junctional adhesion molecule (g3462455) (Mus musculus) | BLAST MOTIFS PFAM SPSCAN HMM |
| 7 | 148 | S26 T106 T4 S36 S40 S142 | | Ig domain G34-R117 Signal peptide M1-S19 | Ig heavy chain variable region (g3170981) (Homo sapiens) | BLAST MOTIFS PFAM SPSCAN HMM |
| 8 | 310 | T11 S303 T137 T236 S300 | N183 | Ig domains: G86-Y144 G181-S241 Transmembrane domains: E275-F295 | ILTlc: myeloid- and lymphoid-specific Ig-like receptor (g1907323) (Homo sapiens) | BLAST MOTIFS PFAM HMM |
| 9 | 236 | S36 S89 T124 S184 T186 S204 T44 Y54 | | Ig domains G38-Q112 S149-V218 D192-E235 Signal peptide M1-C22 | Ig K light chain (g2765423) (Homo sapiens) | BLAST MOTIFS PROFILES BLOCKS PFAM SPSCAN HMM |

Table 2 cont.

| Polypeptide SEQ ID NO: | Amino Acid Residues | Potential Phosphorylation Sites | Potential Glycosylation Sites | Signature Sequences Motifs, and Domains | Homologous Sequences | Analytical Methods and Databases |
|---------------------------|---------------------------|---------------------------------------|-------------------------------------|--|---|---|
| 10 | 237 | S36 S89 T125 S185 T187 S205 T44 | | Ig domains G38-Q112 S150-V219 D193-E236 Signal peptide M1-C22 | Ig κ light chain (g2765423) (Homo sapiens) | BLAST MOTIFS PROFILES SCAN BLOCKS PFAM SPSCAN HMM |
| 11 | 148 | S26 T106 S36 S40 S142 | | Ig domain G34-R117 Signal peptide M1-S19 | Ig heavy chain variable region (g3170891) (Homo sapiens) | BLAST MOTIFS PFAM SPSCAN HMM |
| 12 | 236 | S36 S74 T124 S184 T186 S204 | | Ig domains G38-H112 S149-V218 D192-E235 Signal peptide M1-C22 | Ig κ light chain (g2765423) (Homo sapiens) | BLAST MOTIFS PROFILES SCAN BLOCKS PFAM SPSCAN HMM |
| 13 | 237 | S36 S89 T125 S185 T187 S205 T44 | | Ig domains G38-Q112 S150-V219 D193-E236 Signal peptide M1-C22 | Ig κ light chain (g2765423) (Homo sapiens) | BLAST MOTIFS PROFILES SCAN BLOCKS PFAM SPSCAN HMM |

Table 2 cont.

| Polypeptide SEQ ID NO: | Amino Acid Residues | Potential Phosphorylation Sites | Potential Glycosylation Sites | Signature Sequences Motifs, and Domains | Homologous Sequences | Analytical Methods and Databases |
|---------------------------|---------------------------|---|-------------------------------------|--|---|--|
| 14 | 219 | S26 T47 T157 T181 S183 T4 S36 S40 S146 T190 | | Ig domain G34-R117 Signal peptide M1-S19 | Ig heavy chain variable region (g386807) (Homo sapiens) | BLAST MOTIFS PFAM BLOCKS SPSCAN HMM |
| 15 | 241 | S18 S92 S101 T129 S189 T191 S209 S42 S77 T97 | | Ig Domain: G36-R115 S154-V223 Signal Peptide: M1-A20 | chimeric monoclonal TSH ab, κ chain (g413074) (synthetic construct) | BLAST PFAM, MOTIFS, BLOCKS, SPSCAN, ProfileScan |
| 16 | 507 | T114 T170 T194 S410 S435 S90 T101 S107 S159 T203 S320 T379 T479 | N298 N494 | Ig Domain: S41-T125 N397-V469 Signal Peptide: M1-S26 | SNC73 protein, down regulated protein in colorectal cancer (g3201900) (Homo sapiens) | BLAST PFAM, MOTIFS, BLOCKS, SPSCAN, ProfileScan |
| 17 | 244 | S38 S82 S97 T132 S192 T194 S212 T46 | | Ig Domain: G40-Q120 S157-V226 Signal Peptide: M1-A20 | anti-Entamoeba histolytica Ig κ light chain (g5360673) (Homo sapiens) | BLAST PFAM, MOTIFS, BLOCKS, SPSCAN, ProfileScan |

Table 2 cont.

| Polypeptide SEQ ID NO: | Amino Acid Residues | Potential Phosphorylation Sites | Potential Glycosylation Sites | Signature Sequences Motifs, and Domains | Homologous Sequences | Analytical Methods and Databases |
|---------------------------|---------------------------|---|--|--|--|--|
| 18 | 240 | S20 S149 S228 S78 S215 | | Ig Domain: G39-L116 A155-V223 Signal Peptide: M1-S24 | Ig λ light chain (g2765427) (Homo sapiens) | BLAST PFAM, MOTIFS, BLOCKS, SPSCAN, ProfileScan |
| 19 | 398 | T88 T107 T152 S199 S74 T194 S286 Y216 | N171 N197 N229 N284 N296 N317 N382 | Ig Domain: G45-A118 G252-A318 Signal Peptide: M1-A28 Transmembrane L340-Y360 | Glycoprotein 56 (g1806278) (Rattus norvegicus) | BLAST PFAM, MOTIFS, BLOCKS, SPSCAN, HMMER, ProfileScan |

Table 3

| Nucleotide SEQ ID NO: | Selected Fragment(s) | Tissue Expression (Fraction of Total) | Diseases, Disorders, or Conditions (Fraction of Total) | Vector |
|--------------------------|---------------------------------|---|--|-------------|
| 20 | 356-400 | Reproductive (0.285) Gastrointestinal (0.261) Cardiovascular (0.116) | Cancer (0.598) Inflammation (0.269) | PBLUESCRIPT |
| 21 | 110-154 | Gastrointestinal (0.308) Reproductive (0.215) Hematopoietic/Immune (0.159) | Cancer (0.514) Inflammation (0.411) | pINCY |
| 22 | 346-393 | Hematopoietic/Immune (0.274) Gastrointestinal (0.233) Reproductive (0.178) | Inflammation (0.411) Cancer (0.356) Fetal (0.110) | pINCY |
| 23 | 659-688 | Hematopoietic/Immune (0.407) Nervous (0.148) Cardiovascular (0.111) | Inflammation (0.556) Cancer (0.333) Fetal (0.074) | PSPORT1 |
| 24 | 209-250 | Gastrointestinal (0.270) Reproductive (0.266) Hematopoietic/Immune (0.129) | Cancer (0.571) Inflammation (0.296) Trauma (0.086) | pINCY |
| 25 | 494-538 | Nervous (0.412) Reproductive (0.265) Cardiovascular (0.088) | Cancer (0.471) Inflammation (0.206) Fetal (0.176) | pINCY |
| 26 | 263-307 | Gastrointestinal (0.324) Reproductive (0.250) Hematopoietic/Immune (0.120) | Cancer (0.546) Inflammation (0.370) Trauma (0.065) | pINCY |
| 27 | 327-371 597-641 1029-1073 | Hematopoietic/Immune (0.562) Gastrointestinal (0.188) Musculoskeletal (0.125) | Inflammation (0.625) Cancer (0.250) Fetal (0.250) | pINCY |

Table 3 cont.

| Nucleotide SEQ ID NO: | Selected Fragment(s) | Tissue Expression (Fraction of Total) | Diseases, Disorders, or Conditions (Fraction of Total) | Vector |
|--------------------------|-------------------------|--|--|--------|
| 28 | 149-232 | Reproductive (0.280) Gastrointestinal (0.259) Hematopoietic/Immune (0.126) | Cancer (0.582) Inflammation (0.285) Trauma (0.084) | pINCY |
| 29 | 154-183 | Reproductive (0.274) Gastrointestinal (0.259) Cardiovascular (0.124) | Cancer (0.586) Inflammation (0.263) Trauma (0.094) | pINCY |
| 30 | 245-292 | Gastrointestinal (0.316) Reproductive (0.194) Cardiovascular (0.143) | Cancer (0.571) Inflammation (0.337) Trauma (0.071) | pINCY |
| 31 | 152-187 | Gastrointestinal (0.281) Reproductive (0.255) Hematopoietic/Immune (0.128) | Cancer (0.570) Inflammation (0.294) Trauma (0.085) | pINCY |
| 32 | 360-389 | Reproductive (0.281) Gastrointestinal (0.258) Cardiovascular (0.123) | Cancer (0.585) Inflammation (0.269) Trauma (0.092) | pINCY |
| 33 | 380-430 | Gastrointestinal (0.306) Reproductive (0.218) Cardiovascular (0.137) | Cancer (0.565) Inflammation (0.339) Trauma (0.081) | pINCY |
| 34 | 70-114 355-396 | Gastrointestinal (0.269) Reproductive (0.269) Hematopoietic/Immune (0.130) Cardiovascular (0.121) | Cancer (0.561) Inflammation (0.305) | pINCY |
| 35 | 80-121 177-466 | Reproductive (0.284) Gastrointestinal (0.276) Cardiovascular (0.133) Hematopoietic/Immune (0.116) | Cancer (0.591) Inflammation (0.307) | pINCY |

Table 3 cont.

| Nucleotide SEQ ID NO: | Selected Fragment(s) | Tissue Expression (Fraction of Total) | Diseases, Disorders, or Conditions (Fraction of Total) | Vector |
|--------------------------|--|--|--|--------|
| 36 | 1-32 (5' UTR, possible promoter) 105-149 393-434 | Gastrointestinal (0.272) Reproductive (0.263) Hematopoietic/Immune (0.129) Cardiovascular (0.121) | Cancer (0.567) Inflammation (0.299) | pINCY |
| 37 | 370-414 | Gastrointestinal (0.303) Reproductive (0.244) Cardiovascular (0.129) Hematopoietic/Immune (0.119) | Cancer (0.562) Inflammation (0.308) | pINCY |
| 38 | 2-169 (5'-UTR, possible promoter) 1250-1363 | Reproductive (0.245) Nervous (0.214) Gastrointestinal (0.153) Cardiovascular (0.133) | Cancer (0.408) Inflammation (0.255) | pINCY |

Table 4

| Nucleotide SEQ ID NO: | Library | Library Description |
|--------------------------|-----------|---|
| 20 | SYNORAB01 | The SYNORAB01 cDNA library was constructed using RNA isolated from the synovial membrane tissue of a 68-year-old Caucasian female with rheumatoid arthritis. |
| 21 | THYRN0T08 | The THYRN0T08 cDNA library was constructed using RNA isolated from diseased left thyroid tissue removed from a 13-year-old Caucasian female during a complete thyroidectomy. Pathology indicated lymphocytic thyroiditis. |
| 22 | THYMN0T05 | The THYMN0T05 cDNA library was constructed using RNA isolated from thymus tissue removed from a 3-year-old Hispanic male during thymectomy and closure of a patent ductus arteriosus. Patient history included cardiac catheterization. |
| 23 | PROSTUT03 | The library was constructed using RNA isolated from prostate tumor tissue removed from a 67-year-old Caucasian male during radical prostatectomy and lymph node biopsy. Pathology indicated adenocarcinoma Gleason grade 3+3. Adenofibromatous hyperplasia was present. Patient history included coronary artery disease, stomach ulcer, and osteoarthritis. Family history included congestive heart failure. |
| 24 | LATRTUT02 | The library was constructed using RNA isolated from a myxoma removed from the left atrium of a 43-year-old Caucasian male during annuloplasty. Pathology indicated atrial myxoma. Patient history included pulmonary insufficiency, acute myocardial infarction, atherosclerotic coronary artery disease, hyperlipidemia, and tobacco use. Family history included benign hypertension, acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes. |
| 25 | SKINBIT01 | The library was constructed using RNA isolated from diseased skin tissue of the left lower leg. Patient history included erythema nodosum of the left lower leg. |

Table 4 cont.

| Nucleotide SEQ ID NO: | Library | Library Description |
|--------------------------|-----------|--|
| 26 | BLADTUT07 | The library was constructed using RNA isolated from bladder tumor tissue removed from the anterior bladder wall of a 58-year-old Caucasian male during a radical cystectomy, radical prostatectomy, and gastrectomy. Pathology indicated a grade 3 transitional cell carcinoma in the left lateral bladder. Patient history included acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes. |
| 27 | COLANOT02 | The library was constructed using RNA isolated from diseased ascending colon tissue removed from a 25-year-old Caucasian female during a multiple segmental resection of the large bowel. Pathology indicated moderately to severely active chronic ulcerative colitis, involving the entire colectomy specimen and sparing 2 cm of the attached ileum. Grossly, the specimen showed continuous involvement from the rectum proximally; marked mucosal atrophy and no skip areas were identified. Microscopically, the specimen showed dense, predominantly mucosal inflammation and crypt abscesses. Patient history included benign large bowel neoplasm. Previous surgeries included a polypectomy. |
| 28 | BRSTTUT13 | The library was constructed using RNA isolated from breast tumor tissue removed from the right breast of a 46-year-old Caucasian female during a unilateral extended simple mastectomy with breast reconstruction. Pathology indicated an invasive grade 3 adenocarcinoma, ductal type with apocrine features and greater than 50% intraductal component. Patient history included breast cancer. |
| 29 | COLAUCT01 | The library was constructed using RNA isolated from diseased ascending colon tissue removed from a 74-year-old Caucasian male during a multiple-segment large bowel excision with temporary ileostomy. Pathology indicated inflammatory bowel disease most consistent with chronic ulcerative colitis, characterized by severe acute and chronic mucosal inflammation with erythema, ulceration, and pseudopolyp formation involving the entire colon and rectum. The sigmoid colon had an area of mild stricture formation. One diverticulum with diverticulitis was identified near this zone. |

Table 4 cont.

| Nucleotide SEQ ID NO: | Library | Library Description |
|--------------------------|-----------|---|
| 30 | LUNGNOT31 | The library was constructed using RNA isolated from right middle lobe lung tissue removed from a 63-year-old Caucasian male. Pathology for the associated tumor indicated grade 3 adenocarcinoma. Patient history included an abdominal aortic aneurysm, cardiac dysrhythmia, atherosclerotic coronary artery disease, hiatal hernia, chronic sinusitis, and lupus. Family history included acute myocardial infarction and atherosclerotic coronary artery disease. |
| 31 | BMARNOT03 | The library was constructed using RNA isolated from the left tibial bone marrow tissue of a 16-year-old Caucasian male during a partial left tibial osteotomy with free skin graft. Patient history included an abnormality of the red blood cells. Previous surgeries included bone and bone marrow biopsy, and soft tissue excision. Family history included osteoarthritis. |
| 32 | LUNGUT08 | The library was constructed using RNA isolated from lung tumor tissue removed from a 63-year-old Caucasian male during a right upper lobectomy with fiberoptic bronchoscopy. Pathology indicated a grade 3 adenocarcinoma. Patient history included atherosclerotic coronary artery disease, acute myocardial infarction, rectal cancer, asymptomatic abdominal aortic aneurysm, tobacco abuse, and cardiac dysrhythmia. Family history included congestive heart failure, stomach cancer, lung cancer, type II diabetes, atherosclerotic coronary artery disease, and acute myocardial infarction. |
| 33 | GBLADIT02 | Library was constructed using RNA isolated from diseased gallbladder tissue removed from an 18-year-old Caucasian female during cholecystectomy and incidental appendectomy. Pathology indicated acute and chronic cholecystitis with cholelithiasis. The gallbladder contained multiple fragments of stony material. The appendix showed lymphoid hyperplasia. Previous surgeries included total splenectomy. Family history included benign hypertension, acute myocardial infarction, and atherosclerotic coronary artery disease. |

Table 4 cont.

| Nucleotide SEQ ID NO: | Library | Library Description |
|--------------------------|-----------|--|
| 34 | COLAUCT01 | The COLAUCT01 library was constructed using RNA isolated from diseased ascending colon tissue removed from a 74-year-old Caucasian male during a multiple-segment large bowel excision with temporary ileostomy. Pathology indicated inflammatory bowel disease most consistent with chronic ulcerative colitis, characterized by severe acute and chronic mucosal inflammation with erythema, ulceration, and pseudopolyp formation involving the entire colon and rectum. The sigmoid colon had an area of mild stricture formation. One diverticulum with diverticulitis was identified near this zone. |
| 35 | SINTNOT25 | The SINTNOT25 library was constructed using RNA isolated from small intestine tissue removed from a 13-year-old Caucasian male, who died from a gunshot wound to the head. Family history included diabetes. |
| 36 | SINTNOT25 | The SINTNOT25 library was constructed using RNA isolated from small intestine tissue removed from a 13-year-old Caucasian male, who died from a gunshot wound to the head. Family history included diabetes. |
| 37 | SINTNOT25 | The SINTNOT25 library was constructed using RNA isolated from small intestine tissue removed from a 13-year-old Caucasian male, who died from a gunshot wound to the head. Family history included diabetes. |
| 38 | BRABDIR01 | The BRABDIR01 library was constructed using RNA isolated from diseased cerebellum tissue removed from the brain of a 57-year-old Caucasian male, who died from a cerebrovascular accident. |

Table 5

| Program | Description | Reference | Parameter Threshold |
|-------------------|---|--|---|
| ABI FACTURA | A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences. | Perkin-Elmer Applied Biosystems, Foster City, CA. | |
| ABI/PARACEL FDF | A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences. | Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA. | Mismatch <50% |
| ABI AutoAssembler | A program that assembles nucleic acid sequences. | Perkin-Elmer Applied Biosystems, Foster City, CA. | |
| BLAST | A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx. | Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402. | ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less |
| FASTA | A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch. | Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489. | ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater |
| BLIMPS | A BLOCKS IMPROVED Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions. | Henikoff, S. and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424. | Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less |
| HMMER | An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM. | Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322. | Score=10-50 bits for PFAM hits, depending on individual protein families |

Table 5 (cont.)

| Program | Description | Reference | Parameter Threshold |
|-------------|---|--|--|
| ProfileScan | An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite. | Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221. | Normalized quality score \geq GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1. |
| Phred | A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability. | Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194. | |
| Phrap | A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences. | Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA. | Score= 120 or greater; Match length= 56 or greater |
| Consed | A graphical tool for viewing and editing Phrap assemblies | Gordon, D. et al. (1998) Genome Res. 8:195-202. | |
| SPScan | A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides. | Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439. | Score=3.5 or greater |
| Motifs | A program that searches amino acid sequences for patterns that matched those defined in Prosite. | Bairoch et al. <i>supra</i> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI. | |

What is claimed is:

1. A substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-19 and fragments thereof.

2. A substantially purified variant having at least 90% amino acid sequence identity to the amino acid sequence of claim 1.

3. An isolated and purified polynucleotide encoding the polypeptide of claim 1.

4. An isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide of claim 3.

5. An isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide of claim 3.

6. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 3.

7. A method for detecting a polynucleotide, the method comprising the steps of:

(a) hybridizing the polynucleotide of claim 6 to at least one nucleic acid in a sample, thereby forming a hybridization complex; and

(b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of the polynucleotide in the sample.

8. The method of claim 7 further comprising amplifying the polynucleotide prior to hybridization.

9. An isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:20-38 and fragments thereof.

10. An isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide of claim 9.

11. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 9.

12. An expression vector comprising at least a fragment of the polynucleotide of claim 3.

13. A host cell comprising the expression vector of claim 12.

14. A method for producing a polypeptide, the method comprising the steps of:

- a) culturing the host cell of claim 13 under conditions suitable for the expression of the polypeptide; and
- b) recovering the polypeptide from the host cell culture.

15. A pharmaceutical composition comprising the polypeptide of claim 1 in conjunction with a suitable pharmaceutical carrier.

16. A purified antibody which specifically binds to the polypeptide of claim 1.

17. A purified agonist of the polypeptide of claim 1.

18. A purified antagonist of the polypeptide of claim 1.

19. A method for treating or preventing a disorder associated with decreased expression or activity of IGFAM, the method comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 15.

20. A method for treating or preventing a disorder associated with increased expression or activity of IGFAM, the method comprising administering to a subject in need of such treatment an effective amount of the antagonist of claim 18.

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.

YUE, Henry

TANG, Y. Tom

CORLEY, Neil C.

GUEGLER, Karl J.

GORGONE, Gina A.

BAUGHN, Mariah R.

LU, Dyung Aina M.

LAL, Preeti

HILLMAN, Jennifer L.

YANG, Junming

<120> IMMUNOGLOBULIN SUPERFAMILY PROTEINS

<130> PF-0643 PCT

<140> To Be Assigned

<141> Herewith

<150> 09/195,853; unassigned; 60/113,635; 60/128,194

<151> 1998-11-19; 1998-11-19; 1998-12-22; 1999-04-07

<160> 38

<170> PERL Program

<210> 1

<211> 237

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 079785CD1

<400> 1

| | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Asp | Met | Arg | Val | Pro | Ala | Gln | Leu | Leu | Gly | Leu | Leu | Leu | Leu |
| 1 | | | | 5 | | | | | | 10 | | | | 15 |
| Trp | Leu | Arg | Gly | Ala | Arg | Cys | Asp | Ile | Gln | Met | Thr | Gln | Ser | Pro |
| | | | | 20 | | | | | | 25 | | | | 30 |
| Ser | Ser | Leu | Ser | Ala | Ser | Val | Gly | Asp | Arg | Val | Thr | Ile | Thr | Cys |
| | | | | 35 | | | | | | 40 | | | | 45 |
| Arg | Ala | Gly | Gln | Ser | Ile | Ser | Ser | Tyr | Leu | Asn | Trp | Tyr | Gln | Gln |
| | | | | 50 | | | | | | 55 | | | | 60 |
| Lys | Pro | Gly | Lys | Ala | Pro | Lys | Leu | Leu | Ile | Tyr | Ala | Ala | Ser | Ser |
| | | | | 65 | | | | | | 70 | | | | 75 |
| Leu | Gln | Ser | Gly | Val | Pro | Ser | Arg | Phe | Ser | Gly | Ser | Gly | Ser | Gly |
| | | | | 80 | | | | | | 85 | | | | 90 |
| Thr | Asp | Phe | Thr | Leu | Thr | Ile | Ser | Ser | Leu | Gln | Pro | Glu | Asp | Phe |
| | | | | 95 | | | | | | 100 | | | | 105 |
| Ala | Thr | Tyr | Tyr | Cys | Gln | Gln | Ser | Tyr | Ser | Thr | Pro | Pro | Ile | Thr |
| | | | | 110 | | | | | | 115 | | | | 120 |

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|
| Phe | Gly | Gln | Gly | Thr | Arg | Leu | Glu | Ile | Lys | Arg | Thr | Val | Ala | Ala | |
| | | | | 125 | | | | | 130 | | | | | 135 | |
| Pro | Ser | Val | Phe | Ile | Phe | Pro | Pro | Ser | Asp | Glu | Gln | Leu | Lys | Ser | |
| | | | | 140 | | | | | 145 | | | | | 150 | |
| Gly | Thr | Ala | Ser | Val | Val | Cys | Leu | Leu | Asn | Asn | Phe | Tyr | Pro | Arg | |
| | | | | 155 | | | | | 160 | | | | | 165 | |
| Glu | Ala | Lys | Val | Gln | Trp | Lys | Val | Asp | Asn | Ala | Leu | Gln | Ser | Gly | |
| | | | | 170 | | | | | 175 | | | | | 180 | |
| Asn | Ser | Gln | Glu | Ser | Val | Thr | Glu | Gln | Asp | Ser | Lys | Asp | Ser | Thr | |
| | | | | 185 | | | | | 190 | | | | | 195 | |
| Tyr | Ser | Leu | Ser | Ser | Thr | Leu | Thr | Leu | Ser | Lys | Ala | Asp | Tyr | Glu | |
| | | | | 200 | | | | | 205 | | | | | 210 | |
| Lys | His | Lys | Val | Tyr | Ala | Cys | Glu | Val | Thr | His | Gln | Gly | Leu | Ser | |
| | | | | 215 | | | | | 220 | | | | | 225 | |
| Ser | Pro | Val | Thr | Lys | Ser | Phe | Asn | Arg | Gly | Glu | Cys | | | | |
| | | | | 230 | | | | | 235 | | | | | | |

<210> 2

<211> 537

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 2469025CD1

<400> 2

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|
| Met | Asp | Leu | Leu | His | Lys | Asn | Met | Lys | His | Leu | Trp | Phe | Phe | Leu | |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | |
| Leu | Leu | Val | Ala | Ala | Pro | Arg | Trp | Val | Leu | Ser | Gln | Val | Gln | Leu | |
| | | | | 20 | | | | | 25 | | | | | 30 | |
| Gln | Gln | Trp | Gly | Ala | Gly | Leu | Leu | Lys | Pro | Ser | Glu | Thr | Leu | Ser | |
| | | | | 35 | | | | | 40 | | | | | 45 | |
| Leu | Thr | Cys | Ala | Val | Tyr | Gly | Gly | Ser | Phe | Ser | Gly | Tyr | Tyr | Leu | |
| | | | | 50 | | | | | 55 | | | | | 60 | |
| Ser | Gly | Tyr | Tyr | Trp | Ser | Trp | Ile | Arg | Gln | Pro | Pro | Gly | Lys | Gly | |
| | | | | 65 | | | | | 70 | | | | | 75 | |
| Leu | Glu | Trp | Ile | Gly | Glu | Ile | Asn | His | Ser | Gly | Ser | Thr | Asn | Tyr | |
| | | | | 80 | | | | | 85 | | | | | 90 | |
| Asn | Pro | Ser | Leu | Lys | Ser | Arg | Val | Thr | Ile | Ser | Val | Asp | Thr | Ser | |
| | | | | 95 | | | | | 100 | | | | | 105 | |
| Lys | Asn | Gln | Phe | Ser | Leu | Lys | Leu | Ser | Ser | Val | Thr | Ala | Ala | Asp | |
| | | | | 110 | | | | | 115 | | | | | 120 | |
| Thr | Ala | Val | Tyr | Tyr | Cys | Ala | Arg | Gly | Arg | Ser | Asp | Ser | Ser | Gly | |
| | | | | 125 | | | | | 130 | | | | | 135 | |
| Ser | Pro | Tyr | Gly | Leu | Asp | Tyr | Trp | Gly | Gln | Gly | Thr | Leu | Val | Thr | |
| | | | | 140 | | | | | 145 | | | | | 150 | |
| Val | Ser | Ser | Ala | Pro | Thr | Lys | Ala | Pro | Asp | Val | Phe | Pro | Ile | Ile | |
| | | | | 155 | | | | | 160 | | | | | 165 | |
| Ser | Gly | Cys | Arg | His | Pro | Lys | Asp | Asn | Ser | Pro | Val | Val | Leu | Ala | |
| | | | | 170 | | | | | 175 | | | | | 180 | |
| Cys | Leu | Ile | Thr | Gly | Tyr | His | Pro | Thr | Ser | Val | Thr | Val | Thr | Trp | |
| | | | | 185 | | | | | 190 | | | | | 195 | |
| Tyr | Met | Gly | Thr | Gln | Ser | Gln | Pro | Gln | Arg | Thr | Phe | Pro | Glu | Ile | |

200 205 210
 Gln Arg Arg Asp Ser Tyr Tyr Met Thr Ser Ser Gln Leu Ser Thr
 215 220 225
 Pro Leu Gln Gln Trp Arg Gln Gly Glu Tyr Lys Cys Val Val Gln
 230 235 240
 His Thr Ala Ser Lys Ser Lys Lys Glu Ile Phe Arg Trp Pro Glu
 245 250 255
 Ser Pro Lys Ala Gln Ala Ser Ser Val Pro Thr Ala Gln Pro Gln
 260 265 270
 Ala Glu Gly Ser Leu Ala Lys Ala Thr Thr Ala Pro Ala Thr Thr
 275 280 285
 Arg Asn Thr Gly Arg Gly Gly Glu Glu Lys Lys Glu Lys Glu
 290 295 300
 Lys Glu Glu Gln Glu Glu Arg Glu Thr Lys Thr Pro Glu Cys Pro
 305 310 315
 Ser His Thr Gln Pro Leu Gly Val Tyr Leu Leu Thr Pro Ala Val
 320 325 330
 Gln Asp Leu Trp Leu Arg Asp Lys Ala Thr Phe Thr Cys Phe Val
 335 340 345
 Val Gly Ser Asp Leu Lys Asp Ala His Leu Thr Trp Glu Val Ala
 350 355 360
 Gly Lys Val Pro Thr Gly Gly Val Glu Glu Gly Leu Leu Glu Arg
 365 370 375
 His Ser Asn Gly Ser Gln Ser Gln His Ser Arg Leu Thr Leu Pro
 380 385 390
 Arg Ser Leu Trp Asn Ala Gly Thr Ser Val Thr Cys Thr Leu Asn
 395 400 405
 His Pro Ser Leu Pro Pro Gln Arg Leu Met Ala Leu Arg Glu Pro
 410 415 420
 Ala Ala Gln Ala Pro Val Lys Leu Ser Leu Asn Leu Leu Ala Ser
 425 430 435
 Ser Asp Pro Pro Glu Ala Ala Ser Trp Leu Leu Cys Glu Val Ser
 440 445 450
 Gly Phe Ser Pro Pro Asn Ile Leu Leu Met Trp Leu Glu Asp Gln
 455 460 465
 Arg Glu Val Asn Thr Ser Gly Phe Ala Pro Ala Arg Pro Pro Pro
 470 475 480
 Gln Pro Gly Ser Thr Thr Phe Trp Ala Trp Ser Val Leu Arg Val
 485 490 495
 Pro Ala Pro Pro Ser Pro Gln Pro Ala Thr Tyr Thr Cys Val Val
 500 505 510
 Ser His Glu Asp Ser Arg Thr Leu Leu Asn Ala Ser Arg Ser Leu
 515 520 525
 Glu Val Ser Tyr Val Thr Asp His Gly Pro Met Lys
 530 535

<210> 3

<211> 311

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 2906265CD1

<400> 3
 Met Gly Thr Arg Leu Leu Phe Trp Val Ala Phe Cys Leu Leu Gly
 1 5 10 15
 Ala Asp His Thr Gly Ala Gly Val Ser Gln Ser Pro Ser Asn Lys
 20 25 30
 Val Thr Glu Lys Gly Lys Asp Val Glu Leu Arg Cys Asp Pro Ile
 35 40 45
 Ser Gly His Thr Ala Leu Tyr Trp Tyr Arg Gln Ser Leu Gly Gln
 50 55 60
 Gly Leu Glu Phe Leu Ile Tyr Phe Gln Gly Asn Ser Ala Pro Asp
 65 70 75
 Lys Ser Gly Leu Pro Ser Asp Arg Phe Ser Ala Glu Arg Thr Gly
 80 85 90
 Gly Ser Val Ser Thr Leu Thr Ile Gln Arg Thr Gln Gln Glu Asp
 95 100 105
 Ser Ala Val Tyr Leu Cys Ala Ser Ser Phe Leu Ala Gly Arg Gly
 110 115 120
 Asn Thr Ile Tyr Phe Gly Glu Gly Ser Trp Leu Thr Val Val Glu
 125 130 135
 Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala Val Phe Glu Pro
 140 145 150
 Ser Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr Leu Val Cys
 155 160 165
 Leu Ala Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser Trp Trp
 170 175 180
 Val Asn Gly Lys Glu Val His Ser Gly Val Ser Thr Asp Pro Gln
 185 190 195
 Pro Leu Lys Glu Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu
 200 205 210
 Ser Ser Arg Leu Arg Val Ser Ala Thr Phe Trp Gln Asn Pro Arg
 215 220 225
 Asn His Phe Arg Cys Gln Val Gln Phe Tyr Gly Leu Ser Glu Asn
 230 235 240
 Asp Glu Trp Thr Gln Asp Arg Ala Lys Pro Val Thr Gln Ile Val
 245 250 255
 Ser Ala Glu Ala Trp Gly Arg Ala Asp Cys Gly Phe Thr Ser Val
 260 265 270
 Ser Tyr Gln Gln Gly Val Leu Ser Ala Thr Ile Leu Tyr Glu Ile
 275 280 285
 Leu Leu Gly Lys Ala Thr Leu Tyr Ala Val Leu Val Ser Ala Leu
 290 295 300
 Val Leu Met Ala Met Val Lys Arg Lys Asp Phe
 305 310

<210> 4
 <211> 194
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID NO: 788975CD1

<400> 4

Met Thr Met Arg His Asn Trp Thr Pro Asp Leu Ser Pro Leu Trp
 1 5 10 15
 Val Leu Leu Leu Cys Ala His Val Val Thr Leu Leu Val Arg Ala
 20 25 30
 Thr Pro Val Ser Gln Thr Thr Thr Ala Ala Thr Ala Ser Val Arg
 35 40 45
 Ser Thr Lys Asp Pro Cys Pro Ser Gln Pro Pro Val Phe Pro Ala
 50 55 60
 Ala Lys Gln Cys Pro Ala Leu Glu Val Thr Trp Pro Glu Val Glu
 65 70 75
 Val Pro Leu Asn Gly Thr Leu Ser Leu Ser Cys Val Ala Cys Ser
 80 85 90
 Arg Phe Pro Asn Phe Ser Ile Leu Tyr Trp Leu Gly Asn Gly Ser
 95 100 105
 Phe Ile Glu His Leu Pro Gly Arg Leu Trp Glu Gly Ser Thr Ser
 110 115 120
 Arg Glu Arg Gly Ser Thr Gly Thr Gln Leu Cys Lys Ala Leu Val
 125 130 135
 Leu Glu Gln Leu Thr Pro Ala Leu His Ser Thr Asn Phe Ser Cys
 140 145 150
 Val Leu Val Asp Pro Glu Gln Val Val Gln Arg His Val Val Leu
 155 160 165
 Ala Gln Leu Trp Ala Gly Leu Arg Ala Thr Leu Pro Pro Thr Gln
 170 175 180
 Glu Ala Leu Pro Ser Ser His Ser Ser Pro Gln Gln Gln Gly
 185 190

<210> 5

<211> 236

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 1407148CD1

<400> 5

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu
 1 5 10 15
 Trp Leu Pro Gly Ala Arg Cys Asp Ile Gln Leu Thr Gln Ser Pro
 20 25 30
 Ser Phe Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys
 35 40 45
 Arg Ala Ser Gln Leu Ile Ser Asn His Leu Ala Trp Tyr Gln Gln
 50 55 60
 Lys Pro Gly Arg Ala Pro Lys Leu Leu Val His Ser Ala Ser Ile
 65 70 75
 Leu Gln Ser Gly Val Pro Leu Arg Phe Ser Gly Ser Gly Tyr Gly
 80 85 90
 Thr Glu Phe Thr Leu Thr Val Ala Ser Leu Gln Pro Glu Asp Ser
 95 100 105
 Ala Thr Tyr Tyr Cys Gln Gln Arg Asn Gly Tyr Pro Ile Thr Phe
 110 115 120
 Gly Gln Gly Thr Arg Leu Glu Ile Lys Arg Thr Val Ala Ala Pro

```

      125      130      135
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
      140      145      150
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu
      155      160      165
Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn
      170      175      180
Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr
      185      190      195
Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
      200      205      210
His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
      215      220      225
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
      230      235

```

<210> 6

<211> 310

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 1870848CD1

<400> 6

```

Met Ala Leu Arg Arg Pro Pro Arg Leu Arg Leu Cys Ala Arg Leu
  1      5      10      15
Pro Asp Phe Phe Leu Leu Leu Leu Phe Arg Gly Cys Leu Ile Gly
      20      25      30
Ala Val Asn Leu Lys Ser Ser Asn Arg Thr Pro Val Val Gln Glu
      35      40      45
Phe Glu Ser Val Glu Leu Ser Cys Ile Ile Thr Asp Ser Gln Thr
      50      55      60
Ser Asp Pro Arg Ile Glu Trp Lys Lys Ile Gln Asp Glu Gln Thr
      65      70      75
Thr Tyr Val Phe Phe Asp Asn Lys Ile Gln Gly Asp Leu Ala Gly
      80      85      90
Arg Ala Glu Ile Leu Gly Lys Thr Ser Leu Lys Ile Trp Asn Val
      95      100      105
Thr Arg Arg Asp Ser Ala Leu Tyr Arg Cys Glu Val Val Ala Arg
      110      115      120
Asn Asp Arg Lys Glu Ile Asp Glu Ile Val Ile Glu Leu Thr Val
      125      130      135
Gln Val Lys Pro Val Thr Pro Val Cys Arg Val Pro Lys Ala Val
      140      145      150
Pro Val Gly Lys Met Ala Thr Leu His Cys Gln Glu Ser Glu Gly
      155      160      165
His Pro Arg Pro His Tyr Ser Trp Tyr Arg Asn Asp Val Pro Leu
      170      175      180
Pro Thr Asp Ser Arg Ala Asn Pro Arg Phe Arg Asn Ser Ser Ser
      185      190      195
His Leu Asn Ser Glu Thr Gly Thr Leu Val Phe Thr Ala Val His
      200      205      210

```

Lys Asp Asp Ser Gly Gln Tyr Tyr Cys Ile Ala Ser Asn Asp Ala
 215 220 225
 Gly Ser Ala Arg Cys Glu Glu Gln Glu Met Glu Val Tyr Asp Leu
 230 235 240
 Asn Ile Gly Gly Ile Ile Gly Gly Val Leu Val Val Leu Ala Val
 245 250 255
 Leu Ala Leu Ile Thr Leu Gly Ile Cys Cys Ala Tyr Arg Arg Gly
 260 265 270
 Tyr Phe Ile Asn Asn Lys Gln Asp Gly Glu Ser Tyr Lys Asn Pro
 275 280 285
 Gly Lys Pro Asp Gly Val Asn Tyr Ile Arg Thr Asp Glu Glu Gly
 290 295 300
 Asp Phe Arg His Lys Ser Ser Phe Val Ile
 305 310

<210> 7
 <211> 148
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID NO: 1888468CD1

<400> 7
 Met Asp Trp Thr Trp Arg Ile Leu Phe Leu Val Ala Ala Ala Thr
 1 5 10 15
 Gly Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val
 20 25 30
 Lys Lys Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly
 35 40 45
 Tyr Thr Phe Thr Gly Tyr Tyr Met His Trp Val Arg Gln Ala Pro
 50 55 60
 Gly Gln Gly Leu Glu Trp Met Gly Trp Ile Ser Pro Asn Asn Gly
 65 70 75
 Asp Thr Phe Tyr Ala His Arg Leu Gln Asp Arg Val Thr Leu Thr
 80 85 90
 Thr Asp Thr Ser Ala Thr Thr Gly Tyr Met Glu Leu Arg Ser Leu
 95 100 105
 Thr Ser Asp Asp Thr Ala Ile Tyr Tyr Cys Ala Arg Gly Asp Tyr
 110 115 120
 Gly Asn Ser Leu Asp His Trp Gly Gln Gly Asn Leu Val Thr Val
 125 130 135
 Ser Ser Ala Ser Pro Thr Ser Pro Lys Gly Leu Pro Ala
 140 145

<210> 8
 <211> 310
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature

<223> Incyte ID NO: 2770104CD1

<400> 8

| | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Arg | Arg | Thr | Gln | Pro | Leu | Ser | Val | His | Thr | Gly | Trp | Glu | Gly |
| 1 | | | | 5 | | | | | 10 | | | | | 15 |
| Gly | Glu | Ala | Ile | Ser | Leu | Cys | Val | Ser | Leu | Ser | Arg | Gln | His | Arg |
| | | | | 20 | | | | | 25 | | | | | 30 |
| Gly | Leu | Ile | His | Pro | Gln | Ser | Arg | Ala | Val | Gly | Gly | Asp | Ala | Met |
| | | | | 35 | | | | | 40 | | | | | 45 |
| Thr | Pro | Ile | Val | Thr | Val | Leu | Ile | Cys | Leu | Gly | Leu | Ser | Leu | Gly |
| | | | | 50 | | | | | 55 | | | | | 60 |
| Pro | Arg | Thr | His | Val | Gln | Thr | Gly | Thr | Ile | Pro | Lys | Pro | Thr | Leu |
| | | | | 65 | | | | | 70 | | | | | 75 |
| Trp | Ala | Glu | Pro | Asp | Ser | Val | Ile | Thr | Gln | Gly | Ser | Pro | Val | Thr |
| | | | | 80 | | | | | 85 | | | | | 90 |
| Leu | Ser | Cys | Gln | Gly | Ser | Leu | Glu | Ala | Gln | Glu | Tyr | Arg | Leu | Tyr |
| | | | | 95 | | | | | 100 | | | | | 105 |
| Arg | Glu | Lys | Lys | Ser | Ala | Ser | Trp | Ile | Thr | Arg | Ile | Arg | Pro | Glu |
| | | | | 110 | | | | | 115 | | | | | 120 |
| Leu | Val | Lys | Asn | Gly | Gln | Phe | His | Ile | Pro | Ser | Ile | Thr | Trp | Glu |
| | | | | 125 | | | | | 130 | | | | | 135 |
| His | Thr | Gly | Arg | Tyr | Gly | Cys | Gln | Tyr | Tyr | Ser | Arg | Ala | Arg | Trp |
| | | | | 140 | | | | | 145 | | | | | 150 |
| Ser | Glu | Leu | Ser | Asp | Pro | Leu | Val | Ala | Gly | Asp | Asp | Arg | Ser | Tyr |
| | | | | 155 | | | | | 160 | | | | | 165 |
| Gln | Asn | Pro | Thr | Ser | Gln | Pro | Ser | Pro | Gly | Pro | Val | Val | Thr | Pro |
| | | | | 170 | | | | | 175 | | | | | 180 |
| Gly | Lys | Asn | Val | Thr | Leu | Leu | Cys | Gln | Ser | Arg | Gly | Gln | Phe | His |
| | | | | 185 | | | | | 190 | | | | | 195 |
| Thr | Phe | Leu | Leu | Thr | Lys | Glu | Gly | Ala | Gly | His | Pro | Pro | Leu | His |
| | | | | 200 | | | | | 205 | | | | | 210 |
| Leu | Arg | Ser | Glu | His | Gln | Ala | Gln | Gln | Asn | Gln | Ala | Glu | Phe | Arg |
| | | | | 215 | | | | | 220 | | | | | 225 |
| Met | Gly | Pro | Val | Thr | Ser | Ala | His | Val | Gly | Thr | Tyr | Arg | Cys | Tyr |
| | | | | 230 | | | | | 235 | | | | | 240 |
| Ser | Ser | Leu | Ser | Ser | Asn | Pro | Tyr | Leu | Leu | Ser | Leu | Pro | Ser | Asp |
| | | | | 245 | | | | | 250 | | | | | 255 |
| Pro | Leu | Glu | Leu | Val | Val | Ser | Ala | Ser | Leu | Gly | Gln | His | Pro | Gln |
| | | | | 260 | | | | | 265 | | | | | 270 |
| Asp | Tyr | Thr | Val | Glu | Asn | Leu | Ile | Arg | Met | Gly | Val | Ala | Gly | Leu |
| | | | | 275 | | | | | 280 | | | | | 285 |
| Val | Leu | Val | Val | Leu | Gly | Ile | Leu | Leu | Phe | Glu | Ala | Gln | His | Ser |
| | | | | 290 | | | | | 295 | | | | | 300 |
| Gln | Arg | Ser | Leu | Gln | Asp | Ala | Ala | Gly | Arg | | | | | |
| | | | | 305 | | | | | 310 | | | | | |

<210> 9

<211> 236

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 2851053CD1

```

<400> 10
Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu
  1          5          10          15
Trp Leu Arg Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro
  20          25          30
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys
  35          40          45
Arg Ala Ser Gln Ser Ile Ser Ser Tyr Leu Asn Trp Tyr Gln Gln
  50          55          60
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ala Ala Ser Ser
  65          70          75

```

Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly
 80 85 90
 Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe
 95 100 105
 Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Ser Thr Pro Pro Ile Thr
 110 115 120
 Phe Gly Gln Gly Thr Arg Leu Glu Ile Lys Arg Thr Val Ala Ala
 125 130 135
 Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
 140 145 150
 Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
 155 160 165
 Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly
 170 175 180
 Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
 185 190 195
 Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
 200 205 210
 Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
 215 220 225
 Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 230 235

<210> 11
 <211> 148
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID NO: 3559548CD1

<400> 11
 Met Asp Trp Thr Trp Ser Ile Leu Phe Leu Val Ala Ala Ala Thr
 1 5 10 15
 Gly Ala His Ser Gln Val His Leu Val Gln Ser Gly Ala Glu Val
 20 25 30
 Lys Lys Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly
 35 40 45
 Tyr Thr Phe Thr Ser His Gly Ile Thr Trp Val Arg Gln Ala Pro
 50 55 60
 Gly Gln Gly Leu Glu Trp Met Gly Trp Ile Ser Pro Asn Asn Gly
 65 70 75
 Asp Thr Phe Tyr Ala His Arg Leu Gln Asp Arg Val Thr Leu Thr
 80 85 90
 Thr Asp Thr Ser Ala Thr Thr Gly Tyr Met Glu Leu Arg Ser Leu
 95 100 105
 Thr Ser Asp Asp Thr Ala Ile Tyr Tyr Cys Ala Arg Gly Asp Tyr
 110 115 120
 Gly Asn Ser Leu Asp His Trp Gly Gln Gly Asn Leu Val Thr Val
 125 130 135
 Ser Ser Ala Ser Pro Thr Ser Pro Lys Gly Leu Pro Ala
 140 145

<210> 12
 <211> 236
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID NO: 3872741CD1

<400> 12
 Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu
 1 5 10 15
 Trp Leu Ser Gly Ala Arg Cys Asp Thr Gln Met Thr Gln Ser Pro
 20 25 30
 Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Leu Thr Ile Thr Cys
 35 40 45
 Gln Ala Ser Glu Asp Val Ile Lys Tyr Val Asn Trp Tyr Gln Gln
 50 55 60
 Lys Pro Arg Lys Ala Pro Lys Leu Leu Ile His Asp Ala Ser Asn
 65 70 75
 Leu Glu Thr Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly
 80 85 90
 Thr Leu Phe Thr Phe Thr Ile Ser Asn Leu Gln Pro Glu Asp Val
 95 100 105
 Ala Thr Tyr Tyr Cys Gln His Tyr Ala Ser His Pro Leu Thr Phe
 110 115 120
 Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
 125 130 135
 Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
 140 145 150
 Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu
 155 160 165
 Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn
 170 175 180
 Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr
 185 190 195
 Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
 200 205 210
 His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
 215 220 225
 Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 230 235

<210> 13
 <211> 237
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID NO: 3981428CD1

<400> 13
 Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu

1 5 10 15
 Trp Leu Arg Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro
 20 25 30
 Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Met Thr Cys
 35 40 45
 Arg Ala Ser Gln Ser Ile Ser Thr Tyr Leu Asn Trp Tyr Gln Gln
 50 55 60
 Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ala Ala Ser Ser
 65 70 75
 Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly
 80 85 90
 Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe
 95 100 105
 Ala Thr Tyr Tyr Cys Gln Gln Ser Phe Asn Thr His Met Tyr Thr
 110 115 120
 Phe Gly Gln Gly Thr Arg Leu Glu Met Lys Arg Thr Val Ala Ala
 125 130 135
 Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
 140 145 150
 Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
 155 160 165
 Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly
 170 175 180
 Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
 185 190 195
 Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
 200 205 210
 Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
 215 220 225
 Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 230 235

<210> 14

<211> 219

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 4635039CD1

<400> 14

Met Asp Trp Thr Trp Arg Ile Leu Phe Leu Val Ala Ala Val Thr
 1 5 10 15
 Gly Val His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val
 20 25 30
 Arg Lys Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly
 35 40 45
 Tyr Thr Phe Ser Asp His Tyr Ile His Trp Val Arg Gln Ala Pro
 50 55 60
 Gly Gln Gly Leu Glu Trp Met Gly Trp Ile Asn Pro Asn Ser Gly
 65 70 75
 Gly Ala Arg Tyr Ala Gln Gly Phe Gln Gly Leu Val Thr Met Thr
 80 85 90

Arg Asp Thr Ser Ile Ser Thr Ala Tyr Leu Glu Leu Arg Gly Leu
 95 100 105
 Arg Ser Asp Gly Ser Ala Val Tyr Phe Cys Ala Arg Gln Thr Thr
 110 115 120
 Ser Ser Pro Val Gly Asp Ala Phe Asp Ile Trp Gly Gln Gly Thr
 125 130 135
 Met Val Thr Val Ser Ser Ala Ser Pro Thr Ser Pro Lys Val Phe
 140 145 150
 Pro Leu Ser Leu Cys Ser Thr Gln Pro Asp Gly Asn Val Val Ile
 155 160 165
 Ala Cys Leu Val Gln Gly Phe Phe Pro Gln Glu Pro Leu Ser Val
 170 175 180
 Thr Trp Ser Glu Thr Asp Gln Gly Val Thr Ala Lys Lys Leu Pro
 185 190 195
 Thr Gln Pro Gly Cys Leu Arg Gly Thr Val Asn His Glu Gln Pro
 200 205 210
 Ala Asp Pro Ala Gly Gln Asn Ser Ala
 215

<210> 15

<211> 241

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 3240710CD1

<400> 15

Met Arg Leu Pro Ala Gln Leu Leu Gly Leu Leu Met Leu Trp Ile
 1 5 10 15
 Pro Gly Ser Ser Ala Asp Ile Val Leu Thr Gln Thr Pro Leu Ser
 20 25 30
 Leu Ser Val Thr Pro Gly Gln Pro Ala Ser Ile Ser Cys Lys Ser
 35 40 45
 Ser Glu Ser Leu Leu His Thr Asp Gly Lys Thr Tyr Leu His Trp
 50 55 60
 Phe Val Gln Lys Ala Gly Gln Pro Pro Gln Val Leu Met Tyr Glu
 65 70 75
 Val Ser Asn Arg Phe Ser Gly Val Pro Asp Arg Phe Ser Gly Ser
 80 85 90
 Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile Ser Arg Val Glu Ala
 95 100 105
 Glu Asp Val Arg Ile Tyr Tyr Cys Met Arg Thr Ile Gln Val Pro
 110 115 120
 Pro Thr Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
 125 130 135
 Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
 140 145 150
 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
 155 160 165
 Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala
 170 175 180
 Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser

| | | | | | |
|---|-----|--|-----|--|-----|
| | 185 | | 190 | | 195 |
| Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys | | | | | |
| | 200 | | 205 | | 210 |
| Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His | | | | | |
| | 215 | | 220 | | 225 |
| Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu | | | | | |
| | 230 | | 235 | | 240 |
| Cys | | | | | |

<210> 16

<211> 507

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 4945813CD1

<400> 16

| | | |
|---|-----|---------|
| Met Asp Leu Leu Cys Lys Asn Met Lys His Leu Trp Phe Phe Leu | | |
| 1 | 5 | 10 15 |
| Leu Leu Val Ala Ala Pro Arg Trp Val Leu Ser Gln Leu Gln Leu | | |
| | 20 | 25 30 |
| Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu Thr Leu Ser | | |
| | 35 | 40 45 |
| Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Tyr Asn His | | |
| | 50 | 55 60 |
| Tyr Trp Gly Trp Val Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp | | |
| | 65 | 70 75 |
| Ile Gly Ser Ile Phe Tyr Thr Gly Asn Ser Tyr Tyr Asn Pro Ser | | |
| | 80 | 85 90 |
| Leu Lys Ser Arg Leu Ala Ile Ser Val Asp Thr Ser Lys Ser Gln | | |
| | 95 | 100 105 |
| Leu Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val | | |
| | 110 | 115 120 |
| Tyr Tyr Cys Ala Thr Val Pro Lys Thr Arg Ser Arg Pro Arg Gly | | |
| | 125 | 130 135 |
| Tyr Thr Tyr Gly Pro Phe Asp Phe Trp Gly Gln Gly Thr Leu Val | | |
| | 140 | 145 150 |
| Thr Val Ser Ser Ala Ser Pro Thr Ser Pro Lys Val Phe Pro Leu | | |
| | 155 | 160 165 |
| Ser Leu Cys Ser Thr Gln Pro Asp Gly Asn Val Val Ile Ala Cys | | |
| | 170 | 175 180 |
| Leu Val Gln Gly Phe Phe Pro Gln Glu Pro Leu Ser Val Thr Trp | | |
| | 185 | 190 195 |
| Ser Glu Ser Gly Gln Gly Val Thr Ala Arg Asn Phe Pro Pro Ser | | |
| | 200 | 205 210 |
| Gln Asp Ala Ser Gly Asp Leu Tyr Thr Thr Ser Ser Gln Leu Thr | | |
| | 215 | 220 225 |
| Leu Pro Ala Thr Gln Cys Leu Ala Gly Lys Ser Val Thr Cys His | | |
| | 230 | 235 240 |
| Val Lys His Tyr Thr Asn Pro Ser Gln Asp Val Thr Val Pro Cys | | |
| | 245 | 250 255 |
| Pro Val Pro Ser Thr Pro Pro Thr Pro Ser Pro Ser Thr Pro Pro | | |

260 265 270
 Thr Pro Ser Pro Ser Cys Cys His Pro Arg Leu Ser Leu His Arg
 275 280 285
 Pro Ala Leu Glu Asp Leu Leu Leu Gly Ser Glu Ala Asn Leu Thr
 290 295 300
 Cys Thr Leu Thr Gly Leu Arg Asp Ala Ser Gly Val Thr Phe Thr
 305 310 315
 Trp Thr Pro Ser Ser Gly Lys Ser Ala Val Gln Gly Pro Pro Glu
 320 325 330
 Arg Asp Leu Cys Gly Cys Tyr Ser Val Ser Ser Val Leu Pro Gly
 335 340 345
 Cys Ala Glu Pro Trp Asn His Gly Lys Thr Phe Thr Cys Thr Ala
 350 355 360
 Ala Tyr Pro Glu Ser Lys Thr Pro Leu Thr Ala Thr Leu Ser Lys
 365 370 375
 Ser Gly Asn Thr Phe Arg Pro Glu Val His Leu Leu Pro Pro Pro
 380 385 390
 Ser Glu Glu Leu Ala Leu Asn Glu Leu Val Thr Leu Thr Cys Leu
 395 400 405
 Ala Arg Gly Phe Ser Pro Lys Asp Val Leu Val Arg Trp Leu Gln
 410 415 420
 Gly Ser Gln Glu Leu Pro Arg Glu Lys Tyr Leu Thr Trp Ala Ser
 425 430 435
 Arg Gln Glu Pro Ser Gln Gly Thr Thr Thr Phe Ala Val Thr Ser
 440 445 450
 Ile Leu Arg Val Ala Ala Glu Asp Trp Lys Lys Gly Asp Thr Phe
 455 460 465
 Ser Cys Met Val Gly His Glu Ala Leu Pro Leu Ala Phe Thr Gln
 470 475 480
 Lys Thr Ile Asp Arg Leu Ala Gly Lys Pro Thr His Val Asn Val
 485 490 495
 Ser Val Val Met Ala Glu Val Asp Gly Thr Cys Tyr
 500 505

<210> 17

<211> 244

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 4948957CD1

<400> 17

Met Val Leu Gln Thr Gln Val Phe Ile Ser Leu Leu Leu Trp Ile
 1 5 10 15
 Ser Val Leu Thr Ala Gly Ala Tyr Gly Asp Ile Val Met Thr Gln
 20 25 30
 Ser Pro Asp Ser Leu Ala Val Ser Leu Gly Glu Arg Ala Thr Ile
 35 40 45
 Thr Cys Lys Ser Ser Gln Ser Val Phe Tyr Asn Ser Asn Asn Lys
 50 55 60
 Asn Tyr Leu Val Trp Tyr Gln Gln Arg Pro Gly Gln Pro Pro Lys
 65 70 75

Met Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val Pro Asp
 80 85 90
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
 95 100 105
 Ser Ser Leu Gln Ala Glu Asp Val Ala Leu Tyr Tyr Cys Gln Gln
 110 115 120
 Tyr Phe Thr Thr Pro Tyr Thr Phe Gly Gln Gly Thr Arg Leu Glu
 125 130 135
 Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro
 140 145 150
 Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu
 155 160 165
 Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val
 170 175 180
 Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu
 185 190 195
 Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr
 200 205 210
 Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu
 215 220 225
 Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn
 230 235 240
 Arg Gly Glu Cys

<210> 18

<211> 240

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 4949649CD1

<400> 18

Met Ser Val Pro Thr Met Ala Trp Met Met Leu Leu Leu Gly Leu
 1 5 10 15
 Leu Ala Tyr Gly Ser Gly Val Asp Ser Gln Thr Val Val Thr Gln
 20 25 30
 Glu Pro Ser Leu Ser Val Ser Pro Gly Gly Thr Val Thr Leu Thr
 35 40 45
 Cys Gly Leu Ala Ser Asp Ser Val Ser Thr Asn Phe Phe Pro Thr
 50 55 60
 Trp Tyr Gln Gln Thr Pro Gly Gln Ala Pro Arg Thr Leu Ile Tyr
 65 70 75
 Ser Thr Ser Thr Arg Ser Ser Gly Val Pro Asp Arg Phe Ser Gly
 80 85 90
 Ser Ile Leu Gly Asn Lys Ala Ala Leu Thr Ile Thr Gly Ala Gln
 95 100 105
 Ala Asp Asp Glu Ser Asp Tyr Tyr Cys Ala Leu Tyr Met Gly Ser
 110 115 120
 Gly Ile Ser Val Phe Gly Gly Gly Thr Lys Val Thr Val Leu Gly
 125 130 135
 Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser
 140 145 150

| | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Glu | Leu | Gln | Ala | Asn | Lys | Ala | Thr | Leu | Val | Cys | Leu | Ile | Ser |
| | | | | 155 | | | | | 160 | | | | | 165 |
| Asp | Phe | Tyr | Pro | Gly | Ala | Val | Thr | Val | Ala | Trp | Lys | Ala | Asp | Ser |
| | | | | 170 | | | | | 175 | | | | | 180 |
| Ser | Pro | Val | Lys | Ala | Gly | Val | Glu | Thr | Thr | Thr | Pro | Ser | Lys | Gln |
| | | | | 185 | | | | | 190 | | | | | 195 |
| Ser | Asn | Asn | Lys | Tyr | Ala | Ala | Ser | Ser | Tyr | Leu | Ser | Leu | Thr | Pro |
| | | | | 200 | | | | | 205 | | | | | 210 |
| Glu | Gln | Trp | Lys | Ser | His | Arg | Ser | Tyr | Ser | Cys | Gln | Val | Thr | His |
| | | | | 215 | | | | | 220 | | | | | 225 |
| Glu | Gly | Ser | Thr | Val | Glu | Lys | Thr | Val | Ala | Pro | Thr | Glu | Cys | Ser |
| | | | | 230 | | | | | 235 | | | | | 240 |

<210> 19

<211> 398

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 5500302CD1

<400> 19

| | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Ser | Gly | Ser | Ser | Leu | Pro | Ser | Ala | Leu | Ala | Leu | Ser | Leu | Leu |
| 1 | | | | 5 | | | | | 10 | | | | | 15 |
| Leu | Val | Ser | Gly | Ser | Leu | Leu | Pro | Gly | Pro | Gly | Ala | Ala | Gln | Asn |
| | | | | 20 | | | | | 25 | | | | | 30 |
| Ala | Gly | Phe | Val | Lys | Ser | Pro | Met | Ser | Glu | Thr | Lys | Leu | Thr | Gly |
| | | | | 35 | | | | | 40 | | | | | 45 |
| Asp | Ala | Phe | Glu | Leu | Tyr | Cys | Asp | Val | Val | Gly | Ser | Pro | Thr | Pro |
| | | | | 50 | | | | | 55 | | | | | 60 |
| Glu | Ile | Gln | Trp | Trp | Tyr | Ala | Glu | Val | Asn | Arg | Ala | Glu | Ser | Phe |
| | | | | 65 | | | | | 70 | | | | | 75 |
| Arg | Gln | Leu | Trp | Asp | Gly | Ala | Arg | Lys | Arg | Arg | Val | Thr | Val | Asn |
| | | | | 80 | | | | | 85 | | | | | 90 |
| Thr | Ala | Tyr | Gly | Ser | Asn | Gly | Val | Ser | Val | Leu | Arg | Ile | Thr | Arg |
| | | | | 95 | | | | | 100 | | | | | 105 |
| Leu | Thr | Leu | Glu | Asp | Ser | Gly | Thr | Tyr | Glu | Cys | Arg | Ala | Ser | Asn |
| | | | | 110 | | | | | 115 | | | | | 120 |
| Asp | Pro | Lys | Arg | Asn | Asp | Leu | Arg | Gln | Asn | Pro | Ser | Ile | Thr | Trp |
| | | | | 125 | | | | | 130 | | | | | 135 |
| Ile | Arg | Ala | Gln | Ala | Thr | Ile | Ser | Val | Leu | Gln | Lys | Pro | Arg | Ile |
| | | | | 140 | | | | | 145 | | | | | 150 |
| Val | Thr | Ser | Glu | Glu | Val | Ile | Ile | Arg | Asp | Ser | Pro | Val | Leu | Pro |
| | | | | 155 | | | | | 160 | | | | | 165 |
| Val | Thr | Leu | Gln | Cys | Asn | Leu | Thr | Ser | Ser | Ser | His | Thr | Leu | Thr |
| | | | | 170 | | | | | 175 | | | | | 180 |
| Tyr | Ser | Tyr | Trp | Thr | Lys | Asn | Gly | Val | Glu | Leu | Ser | Ala | Thr | Arg |
| | | | | 185 | | | | | 190 | | | | | 195 |
| Lys | Asn | Ala | Ser | Asn | Met | Glu | Tyr | Arg | Ile | Asn | Lys | Pro | Arg | Ala |
| | | | | 200 | | | | | 205 | | | | | 210 |
| Glu | Asp | Ser | Gly | Glu | Tyr | His | Cys | Val | Tyr | His | Phe | Val | Ser | Ala |
| | | | | 215 | | | | | 220 | | | | | 225 |
| Pro | Lys | Ala | Asn | Ala | Thr | Ile | Glu | Val | Lys | Ala | Ala | Pro | Asp | Ile |

| | | | | | |
|---|-----|--|-----|--|-----|
| | 230 | | 235 | | 240 |
| Thr Gly His Lys Arg Ser Glu Asn Lys Asn Glu Gly Gln Asp Ala | | | | | |
| | 245 | | 250 | | 255 |
| Thr Met Tyr Cys Lys Ser Val Gly Tyr Pro His Pro Asp Trp Ile | | | | | |
| | 260 | | 265 | | 270 |
| Trp Arg Lys Lys Glu Asn Gly Met Pro Met Asp Ile Val Asn Thr | | | | | |
| | 275 | | 280 | | 285 |
| Ser Gly Arg Phe Phe Ile Ile Asn Lys Glu Asn Tyr Thr Glu Leu | | | | | |
| | 290 | | 295 | | 300 |
| Asn Ile Val Asn Leu Gln Ile Thr Glu Asp Pro Gly Glu Tyr Glu | | | | | |
| | 305 | | 310 | | 315 |
| Cys Asn Ala Thr Asn Ala Ile Gly Ser Ala Ser Val Val Thr Val | | | | | |
| | 320 | | 325 | | 330 |
| Leu Arg Val Arg Ser His Leu Ala Pro Leu Trp Pro Phe Leu Gly | | | | | |
| | 335 | | 340 | | 345 |
| Ile Leu Ala Glu Ile Ile Ile Leu Val Val Ile Ile Val Val Tyr | | | | | |
| | 350 | | 355 | | 360 |
| Glu Lys Arg Lys Arg Pro Asp Glu Val Pro Asp Asp Asp Glu Pro | | | | | |
| | 365 | | 370 | | 375 |
| Ala Gly Pro Met Lys Thr Asn Ser Thr Asn Asn His Lys Asp Lys | | | | | |
| | 380 | | 385 | | 390 |
| Asn Leu Arg Gln Arg Asn Thr Asn | | | | | |
| | 395 | | | | |

<210> 20

<211> 917

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 079785CB1

<400> 20

```

gtcaggacac agcatggaca tgaggggtccc cgctcagctc ctgggggctcc tgctactctg 60
gctccgaggt gccagatgtg acatccagat gaccagctct ccacccctccc tgtctgcatc 120
tgtaggagac agagtcacca tcacttgccg ggcaggctcag agcattagca gctattttaa 180
ttggtatcag cagaaaccag ggaaagcccc taagctcctg atctatgctg catccagttt 240
gcaaagtggg gtcccatcaa gggttcagtgg cagtggatct gggacagatt tcactctcac 300
catcagcagt ctgcaacctg aagattttgc aacttactac tgtcaacaga gttacagtac 360
ccctccgatc accttcggcc aagggacacg actggagatt aaacgaactg tggctgcacc 420
atctgtcttc atcttcccgc catctgatga gcagttgaaa tctggaactg cctctgttgt 480
gtgcctgctg aataacttct atcccagaga ggccaaagta cagtggaagg tggataacgc 540
cctccaatcg ggtaactccc aggagagtgt cacagagcag gacagcaagg acagcaccta 600
cagcctcagc agcaccctga cgctgagcaa agcagactac gagaaacaca aagtctacgc 660
ctgcgaagtc acccatcagg gcctgagctc gcccgtcaca aagagcttca acaggggaga 720
gtgttagagg gagaagtgcc cccacctgct cctcagttcc agcctgaccc cctcccatcc 780
tttggcctct gacctttttt ccacagggga cctaccctta ttgcggctct ccagctcatc 840
tttcacctca cccccctcct cctccttggc tttaattatg ctaatgttgg aggagaaatga 900
ataaataaag tgatcga 917

```

<210> 21

<211> 1746

<212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID NO: 2469025CB1

<400> 21
 ctgctcacat gggaaatact ttctgagagt catggacctc ctgcacaaga acatgaaaca 60
 cctgtgggttc ttctctctcc tgggtggcagc tcccagatgg gtcctgtccc aggtgcagct 120
 acagcagtggtg ggcgaggagc tggtgaagcc ttccggagacc ctgtccctca cctgcgctgt 180
 ctatgggtggg tccttcagtg gttactactt aagtgggttac tactggagct ggatccgcca 240
 gccccagggg aaggggctgg agtggattgg ggaaatcaat catagtggaa gcaccaacta 300
 caaccgctcc ctcaagagtc gagtcacat atcagtagac acgtccaaga accagttctc 360
 cctgaagctg agctctgtga ccgcccggga cagggtgtg tattactgtg cgagaggcag 420
 gagtgatagt agtgggtccc catatggact tgactactgg ggccagggaa ccctgggtcac 480
 cgtctctctca gcaccacca aggtctccga tgtgttcccc atcatatcag ggtgcagaca 540
 cccaaaggat aacagccctg tggctcctggc atgcttgata actgggtacc acccaacgtc 600
 cgtgactgtc acctggtaca tggggacaca gagccagccc cagagaacct tccctgagat 660
 acaaagacgg gacagctact acatgacaag cagccagctc tccaccccc tccagcagtg 720
 gcgccaaggc gactacaaat gcgtgggtcca gcacaccgcc agcaagagta agaaggagat 780
 ctctccgctgg ccagagtctc caaaggcaca ggctctctca gtgccactg cacaaccca 840
 agcagagggg agcctcgcca aggcaaccac agccccagcc accaccgta acacaggaag 900
 aggaggagaa gagaagaaga aggagaagga gaaagaggaa caagaagaga gagagacaaa 960
 gacaccagag tgtccgagcc acaccagcc tcttggcgct tacctgctaa cccctgcagt 1020
 gcaggacctg tggctccggg acaaagccac cttcacctgc ttcgtgggtg gcagtgaact 1080
 gaaggatgct cacctgacct gggagggtggc tgggaaggtc cccacagggg gcgtggagga 1140
 agggctgtct gagcggcaca gcaacggctc ccagagccag cacagccgtc tgacctgcc 1200
 caggtccttg tggaaagcgg ggacctccgt cacctgcaca ctgaaccatc ccagcctccc 1260
 accccagagg ttgatggcgc tgagagaacc cgctgcgcag gcaccctca agctttccct 1320
 gaacctgctg gcctcgtctg acctcccga ggcgccctcg tggctcctgt gtgaggtgtc 1380
 tggcttctcg ccccccaaca tcctcctgat gtggctggag gaccagcgtg aggtgaacac 1440
 ttctgggttt gccccgcac gccccctcc acagcccggg agcaccagt tctgggcctg 1500
 gagtgtgctg cgtgtcccag ccccgcccag cctcagcca gccacctaca cgtgtgtggt 1560
 cagccacgag gactcccgga ctctgctcaa cgccagccgg agcctagaag tcagctatgt 1620
 aacagacat ggccccatga aatgatccc gaccagatcc gtccacacc gccactcagc 1680
 agctctggcc gagctcacag tacaaccaca ataaactctt gttgaatgaa ctctaaaaaa 1740
 aaaaaa 1746

<210> 22
 <211> 1160
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID NO: 2906265CB1

<400> 22
 gccatgggca ccaggctcct cttctgggtg gccttctgtc tcctgggggc agatcacaca 60
 ggagctggag tctcccagtc ccccagtaac aaggtcacag agaagggaaa ggatgtagag 120
 ctccaggtgtg atccaatttc aggtcatact gccctttact ggtaccgaca gagcctgggg 180
 cagggccttg agtttttaat ttacttccaa ggcaacagt caccagacaa atcagggctg 240
 cccagtgatc gcttctctgc agagaggact gggggatccg tctccactct gacgatccag 300

```

cgcacacagc aggaggactc ggccgtgtat ctctgtgcc aagcctttct tgcagggagg 360
ggaaacacca tatattttgg agaggaagtg tggctcactg ttgtagagga cctgaacaag 420
gtgttccccc cagaggtcgc tgtgtttgag ccatcagaag cagagatctc ccacacccaa 480
aaggccacac tgggtgtgcct ggccacaggg ttcttccctg accacgtgga gctgagctgg 540
tgggtgaatg ggaaggaggt gcacagtggg gtcagcacgg acccgagcc cctcaaggag 600
cagcccgcct tcaatgactc cagatactgc ctgagcagcc gcctgagggg ctcgccacc 660
ttctggcaga acccccga ccacttccgc tgtcaagtcc agttctacgg gctctcggag 720
aatgacgagt ggacccagga tagggccaaa cccgtcacc agatcgtcag cgcagaggcc 780
tggggtagag cagactgtgg ctttacctcg gtgtcctacc agcaaggggt cctgtctgcc 840
accatcctct atgagatcct gctaggggag gccaccctgt atgtgtgct ggtcagcgcc 900
cttgtgttga tggccatggg caagagaaag gatttctgaa ggagccctg gaagtggagt 960
taggagcttc taaccgtc tgggttcaat acacattctt cttttgccag cgcttctgaa 1020
gagctgctct cacctctctg catcccaata gatcccccc tatgtgcatg cacacctgca 1080
cactcacggc tgaaatctcc ctaaccagg gggaccttag catgcctaag tgactaaacc 1140
aaataaaaat gttctgggtca                                     1160

```

<210> 23

<211> 1356

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 788975CB1

<400> 23

```

gctgaagaga gcactgcctc cctgtgtgac tggagaagag gacgttgtca cagataaaga 60
gccaggctca ccagctcctg acgcatgcat catgaccatg agacacaact ggacaccaga 120
cctcagccct ttgtgggtcc tgcctcctgt tgcccacgtc gtcactctcc tgggtcagagc 180
cacacctgtc tcgcagacca ccacagctgc cactgcctca gttagaagca caaaggaccc 240
ctgccccctc cagccccag tgttcccagc agctaagcag tgtccagcat tggaaagtgc 300
ctggccagag ctggaagtgc cactgaatgg aacgctgagc ttatcctgtg tggcctgcag 360
ccgttcccc aacttcagca tcctctactg gctgggcaat ggttcttca ttgagcacct 420
cccaggccga ctgtgggagg ggagcaccag ccgggaacgt gggagcacag gtacgcagct 480
gtgcaaggcc ttggtgctgg agcagctgac cctgcctctg cacagcacca acttctcctg 540
tgtgtctctg gaccctgaac aggttgtcca gcgtcacgtc gtcttgccc agctctgggc 600
tgggctgagg gcaaccttgc cccccaacca agaagccctg cctccagcc acagcagctc 660
acagcagcag ggttaagact cagcacaggg ccagcagcag cacaaccttg accagagctt 720
gggtcctacc tgtctacctg gactgaacag tccctgactg cctgtaggct gcgtggatgc 780
gcaacacacc cctccttct ctgctttggg tcccttctct caccaaattc aaactccatt 840
cccacctacc tagaaaatca cagcctcctt ataatgcctc ctctcctgc cattctctct 900
ccacctatcc attagccttc ctaacgtcct actcctcaca ctgctctact gctcagaaac 960
caccaagact gttgatgcct tagccttgca ctccaggggc ctacctgcat tccccacatg 1020
actttctgga agcctcccaa ctattcttgc ttttccaga cagctccac tcccatgtct 1080
ctgctcattt agtccctct tcctcacgc cccagcaggg gaacgctcaa gcctgggtga 1140
aatgctgcct ctacagtga gtcatcctct ttcatgctct gccgcattct gcagacttcc 1200
tatcttctgt ctgtatgttt ttttttttcc ccttccactc taatggactg ttccagggaa 1260
gggatggggg cagcagctgc ttcggatcca cactgtatct gtgtcatccc cacatgggtc 1320
ctcataaagg attattcaat ggaggcaaaa aaaaaa                                     1356

```

<210> 24

<211> 916

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 1407148CB1

<400> 24

```

gtcaggacac agcatggaca tgaggggtccc cgctcagctc ctgggggtccc tgctgctctg 60
gtcctccaggt gccagatgtg acatccagtt gaccaggtct ccatccttcc tgtctgcgtc 120
tgtgggagac agagtcacca tcacttgccg ggccagtcag ctcatagta atcatttagc 180
ctggtatcag caaaagccag ggagagcccc taaactcctg gtccatagtg catccattct 240
gcaaagtggg gtcccattaa gattcagcgg cagtggatac gggacagagt tcactctcac 300
agtgcgcagc ctgcagcctg aggattccgc aacttattac tgtcaacagc gcaacggtta 360
tccgatcacc ttcccgccat ctgatgagca gttgaaatct ggaactgcct ctggtgtgtg 420
tgtcttcac ttcccgccat ctgatgagca gttgaaatct ggaactgcct ctggtgtgtg 480
cctgctgaat aacttctatc ccagagaggc caaagtacag tgggaagggtg ataacgccct 540
ccaatcgggt aactcccagg agagtgtcac agagcaggac agcaaggaca gcacctacag 600
cctcagcagc accctgacgc tgagcaaagc agactacgag aaacacaaag tctacgcctg 660
cgaagtcacc catcagggcc tgagctcgcc cgtcacaaag agcttcaaca ggggagagtg 720
ttagagggag aagtgcctcc accctgctct cagttccagc ctgacccctt cccatccttt 780
ggcctctgac cctttttcca caggggacct acccctattg cggtcctcca gctcatcttt 840
cacctacccc cctcctcctt ccttggtctt aattatgcta atgttgaggg agaataaata 900
aataaagtga atcttc 916

```

<210> 25

<211> 1956

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 1870848CB1

<400> 25

```

cggggactac aagccgcgcc gcgctgccgc tggccctca gcaaccctcg acatggcgct 60
gaggcggcca ccgcgactcc ggcctctgcgc tcggctgcct gacttcttcc tgcctgctgt 120
tttcaggggc tgctgatag gggtgtgaaa tctcaaatcc agcaatcgaa cccagtggt 180
acaggaattt gaaagtgtgg aactgtcttg catcattacg gattgcgaga caagtgacct 240
caggatcgag tggagaaaaa ttcaagatga acaaccaca tatgtgtttt ttgacaacaa 300
aattcaggga gacttggcgg gtctgtgcaga aatactgggg aagacatccc tgaagatctg 360
gaatgtgaca cggagagact cagcccttta tcgctgtgag gtctgtgtc gaaatgacct 420
caaggaaatt gatgagattg tgatcgagtt aactgtgcaa gtgaagccag tgaccctgt 480
ctgtagagtg ccgaaggctg taccagtagg caagatggca acactgcact gccaggagag 540
tgagggccac ccccgccctc actacagctg gtatcgcaat gatgtaccac tgcccacgga 600
ttccagagcc aatcccagat ttgcgaattc ttcttcccac ttaactctg aaacaggcac 660
tttggtgttc actgctgttc acaaggacga ctctgggcag tactactgca ttgcttccaa 720
tgacgcaggg ctgcctcagg gtgaggagca ggagatggaa gtctatgacc tgaacattgg 780
cggaattatt gggggggttc tgggtgtcct tgctgtactg gccctgatca cgttgggcat 840
ctgctgtgca tacagacgtg gctacttcat caacaataaa caggatggag aaagttaaaa 900
gaaccaggga aaaccagatg gagttaacta catccgcact gacgaggagg gcgacttcag 960
acacaagtca tcgtttgtga tctgagacct gcggtgtggc tgagagcgca cagagcgcac 1020
gtgcacatac ctctgctaga aactcctgtc aaggcagcga gagctgatgc actcggacag 1080
agctagacac tcattcagaa gcttttctgt ttggccaaag ttgaccata ctcttcttac 1140
tctaacaagc cacatgaata gaagaatttt cctcaagatg gaccggtaa atataaccac 1200

```

```

aaggaagcga aactgggtgc gttcactgag ttgggttcct aatctgtttc tggcctgatt 1260
cccgcatgaa tattaggggtg atcttaaaga gtttgctcac gtaaacgccc gtgctgggccc 1320
ctgtgaagcc agcatgttca ccactggtcg ttcagcagcc acgacagcac catgtgagat 1380
ggcgaggtgg ctggacagca ccagcagcgc atcccggcgg gaaccagaa aaggcttctt 1440
acacagcagc cttacttcat cggcccacag acaccaccgc agtttcttct taaaggctct 1500
gctgatcggg gttgcagtgt ccattgtgga gaagcttttt ggatcagcat tttgtaaaaa 1560
caacaaaaat caggaaggta aattggttgc tggaagaggg atcttgccctg aggaaccctg 1620
cttgtccaac aggggtgtcag gatttaagga aaaccttcgt cttaggctaa gtctgaaatg 1680
gtactgaaat atgcttttct atgggtcttg tttattttat aaaattttac atctaaattt 1740
ttgctaagga tgtattttga ttattgaaaa gaaaattttct atttaaactg taaatatatt 1800
gtcatacaat gttaaataac ctattttttt taaaaaagtt caacttaagg tagaagttcc 1860
aagctactag tgataaaatg ggaaatatca ataattaaga gtattttacc caaggaaatc 1920
tcctcagggg agtttactgt gatggctctt tcccgc 1956

```

<210> 26

<211> 589

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 1888468CB1

<400> 26

```

cagcaaccac atctgtcctc tagagaatcc cctgagagct ccgttcctca ccatggactg 60
gacctggagg atcctcttct tgggtggcagc agccacagga gccactccc aggtgcagct 120
ggtgcagtct ggggctgagg tgaagaagcc tggggcctca gtgaaggctc cctgcaaggc 180
ttctggatac accttcaccg gctactatat gcactgggtg cgacaggccc ctggacaagg 240
gcttgagtgg atgggggtgga tcagccctaa caatggtgac acattctatg cacacagact 300
ccaggacaga gtcaccttga ccacagacac atcggcgacc acaggctaca tggagctgag 360
gagcctgaca tctgacgaca cggccatata ttattgcgag agaggggact acggtaactc 420
tcttgaccac tggggccagg gaaacctggt cactgtctcc tcagcatccc cgaccagccc 480
caaaggctct cccgcttgag cctctgcaag caccagccc agatggggaa ccgtggtcaa 540
tcgcctgccc tgggtccagg ggctttcttc cccaaggagg ccaattaat 589

```

<210> 27

<211> 1388

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 2770104CB1

<400> 27

```

caggaaggag cagacacaag aagcaccagt tctatttgcg gctacatccc ggctctcggt 60
gagacgagga gaaaccagac agacagtggc tgggggtcag gaaagacccc attacagtct 120
gaaatgtctg cagagggccc agtgcatgcn cccacctcag ctctaaaaga atgagagtca 180
ggctctctgg agggcagttc cgcttcttgt gtggctgcag atgacaacac cccatgagaa 240
ggaccagccc tctgagtgtc cacacagggg gggaaggagg ggaggctatt tctctctgtg 300
tgtctctgtc ccgccagcac cgagggtcga tccatccgca gagcagggca gtgggaggag 360
acgccatgac ccccatcgtc acagtccctga tctgtctcgg gctgagctct ggccccagga 420
cccacgtgca gacaggggacc atcccgaagc ccacctgtg ggctgagcca gactctgtga 480

```

```

tcacccaggg gagtcccgtc accctcagtt gtcaggggag ccttgaagcc caggagtacc 540
gtctatatag ggagaaaaaa tcagcatctt ggattacacg gatacgacca gagcttgtga 600
agaacggcca gttccacatc ccatecatca cctgggaaca cacagggcga tatggctgtc 660
agtattacag ccgcgctcgg tggctctgagc tcagtgaacc cctgggttgct ggtgatgaca 720
ggagctacca aaaccaacc tctcagccca gcccaggccc tgtggtgacc ccaggaaaga 780
acgtgaccct gctgtgtcag tcacgggggc agttccacac tttccttctg accaaggagg 840
gggcaggcca tccccactg catctgagat cagagcacca agctcagcag aaccaggctg 900
aattccgcat gggctcctgtg acctcagccc acgtggggac ctacagatgc tacagctcac 960
tcagctccaa cccctacctg ctgtctctcc ccagtgaacc cctggagctc gtggtctcag 1020
catccctagg ccaacacccc caggattaca cagtggagaa tctcatccgc atgggtgtgg 1080
ctggcttggc cctggtgggtc ctcggtgattc tgctatttga ggctcagcac agccagagaa 1140
gcctacaaga tgcagccggg aggtgaacag cagagaggac aatgcacctc tcagcgtggc 1200
ggagcctcag ggacagatct gatgatccca ggaggctctg gaggacaatc taggacctac 1260
attatctgga ctgtatgtg gtcatctcta gagacagcaa tcaatatttg agtgtaagga 1320
aactgtctgg ggtgattcct agaagatcat taaactgtgg tacatttttt tgtctatgaa 1380
aaaaaaaa                                     1388

```

<210> 28

<211> 817

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 2851053CB1

<400> 28

```

cagcaacagc atggacatga gagtcctcgc tcagctcctg gggctcctgc tgctctgttt 60
cccagggtgc agatgtgaca tccagatgac ccagtctcca tcctcactgt ctgcatctgt 120
tgagacaga gtcaccatca cttgtcgggc gagtcaggac attagcaatt atttagcctg 180
gtttcagcag aaaccaggga cagcccctaa gtccctgatc tatgatacat ccagtttgca 240
aagtgggggtc ccatcaaagt tcagcggcag tggatctggg acagatttca ctctcaccat 300
caacagcctg cagcctgaag attttgcaac ttattactgc caacagcatc atagttatcc 360
tcttactttc ggcggaggga ccaagggtgga gatcaaacga actgtggctg caccatctgt 420
cttcatcttc ccgccatctg atgagcagtt gaaatctgga actgcctctg ttgtgtgcct 480
gctgaataac ttctatccca gagaggccaa agtacagtgg aaggtggata acgcccctca 540
atcgggtaac tcccaggaga gtgtcacaga gcaggacagc aaggacagca cctacagcct 600
cagcagcacc ctgacgctga gcaaaagcaga ctacgagaaa ccaaagtctc acgcctgcga 660
agtcacccat cagggcctga gctcgccctg cacaaagagc ttcaacaggg gagagtgtta 720
gagggagaag tgtccccacc tgctcctcag ttcagcctga cccctctcca tcctttgggc 780
tctgaccctt tttccacagg gggactaccc ctattggg                                     817

```

<210> 29

<211> 936

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 3238787CB1

<400> 29

```

tctcagtcag gacacagcat ggacatgagg gtccccgctc agctcctggg gctcctgcta 60

```

```

ctctggctcc gaggtgccag atgtgacatc cagatgaccc agtctccatc ctccctgtct 120
gcatctgtag gagacagagt caccatcact tgccgggcaa gtcagagcat tagcagctat 180
ttaaattggt atcagcagaa accagggaaa gccctaagc tcttgatcta tgctgcatcc 240
agtttgcaaa gtgggggtccc atcaagggtc agtggcagtg gatctgggac agatttcact 300
ctcaccatca gcagtctgca acctgaagat ttgcaactt actactgtca acagagttac 360
agtacccctc cgatcacctt cggccaaggg acacgactgg agattaaacg aactgtggct 420
gcaccatctg tcttcatctt cccgccatct gatgagcagt tgaaatctgg aactgcctct 480
gttgtgtgcc tgctgaataa cttctatccc agagaggcca aagtacagtg gaagggtgat 540
aacgccctcc aatcggttaa ctcccaggag agtgtcacag agcaggacag caaggacagc 600
acctacagcc tcagcagcac cctgacgctg agcaaagcag actacgagaa acacaaagtc 660
tacgctgcg aagtcaccca tcagggcctg agctcgcccg tcacaaagag cttcaacagg 720
ggagagtgtt agagggagaa gtgccccac ctgctcctca gttccagcct gacccctcc 780
catcctttgg cctctgaccc tttttccaca ggggacctac ccttattgcg gtcctccagc 840
tcatctttca cctcaccccc ctctcctcc ttggctttaa ttatgctaata gttggaggag 900
aatgaataaaa taaagtgaat ctttgcaaaa aaaaaa 936

```

<210> 30

<211> 571

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 3559548CB1

<400> 30

```

cttgggagaa tcccctagat cacagctcct caccatggac tggacctgga gcatcctttt 60
cttgggtggca gcagcaacag gtgcccactc ccagggttcac ctggtacagt ctggagctga 120
ggtgaagaag cctggggcct cagtgaaggc ctcttgcaag gcttctggtt acacctttac 180
cagtcatggt atcacctggg tgoggcaggc ccttggaaca gggcttgagt ggatgggggtg 240
gatcagccct aacaatggtg acacattcta tgcacacaga ctccaggaca gagtcacctt 300
gaccacagac acatcggcga ccacaggcta catggagctg aggagcctga catctgacga 360
cacggccata tattattgcg cgagagggga ctacggtaac tctcttgacc actggggcca 420
gggaaacctg gtcactgtct cctcagcatc cccgaccagc cccaaaggct tccccgcttg 480
agcctctgca agcaccagc ccagatgggg aaccgtgggc aatcgctgct cctgggtcca 540
ggggctttct tccccaggg agccaattaa t 571

```

<210> 31

<211> 890

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 3872741CB1

<400> 31

```

gtcccaacca ggacacagca tggacatgag ggtccctgct cagctcctgg ggctcctgct 60
gctctggctc tcagggtgcca gatgtgacac ccagatgacc cagtctccat cctccctgtc 120
tgcactgtgc ggagacagac tcaccatcac ttgccaggcg agtgaggacg tcatcaata 180
tgtgaattgg tatcaacaaa aacctaggaa agcccctaaa ctctgatcc acgatgcatc 240
caatttgga acaggggtcc catcaagggt cagtggaaagt ggatctggga cactttttac 300
tttcaccatc agcaacctgc agcctgaaga tgttgcaaca tattactgtc agcactatgc 360

```

```

tagtcatccg ctcactttcg gcggagggac caaggtggag atcaaacgaa ctgtggctgc 420
accatctgtc ttcactttcc cgccatctga tgagcagttg aaatctggaa ctgcctctgt 480
tgtgtgcctg ctgaataact tctatcccag agaggccaaa gtacagtggg aggtggataa 540
cgccctccaa tcgggtaact cccaggagag tgtcacagag caggacagca aggacagcac 600
ctacagcctc agcagcacc cagcgtgag caaagcagac tacgagaaac acaaagtcta 660
cgctgcgaa gtcacccatc agggcctgag ctgcgccgtc acaaagagct tcaacagggg 720
agagtgttag agggagaagt tgccccacct gctcctcagt tccagcctga cccctccca 780
tcctttggcc tctgagcctt tttccacaag gggactaccc ctattgcggg tctccagctc 840
atctttcacc tcaccccgct cttctgcttg gctttaatta tgctatgttt 890

```

<210> 32

<211> 928

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 3981428CB1

<400> 32

```

ggagtcagtc tcagtcagga cacagcatgg acatgagggc ccccgctcag ctctgggggc 60
tcctgtact ctggctccga ggtgccagat gtgacatcca gatgaccag tctccatcct 120
ccctgtctgc atctgttaga gacagagtca ccatgacttg ccgggcaagt cagagcatta 180
gcacctatct aaattggtat cagcagaaac cagggaagc ccctaagctc ttgatctatg 240
ctgcatccag tttgcaaagt ggggtcccat caagggttcag tggcagtggg tctgggacag 300
atttcactct caccatcagc agtctgcaac ctgaagattt tgcaacttac tactgtcaac 360
agagtttcaa taccacatg tacacttttg gccaggggac acgactggag atgaaacgaa 420
ctgtggctgc accatctgtc ttcactttcc cgccatctga tgagcagttg aaatctggaa 480
ctgcctctgt tgtgtgcctg ctgaataact tctatcccag agaggccaaa gtacagtggg 540
aggtggataa cgccctccaa tcgggtaact cccaggagag tgtcacagag caggacagca 600
aggacagcac ctacagcctc agcagcacc tgacgtgag caaagcagac tacgagaaac 660
acaaagtcta cgctgcgaa gtcacccatc agggcctgag ctgcgccgtc acaaagagct 720
tcaacagggg agagtgttag agggagaagt gccccacct gctcctcagt tccagcctga 780
ccccctccca tcctttggcc tctgacctt tttccacagg ggacctaccc ctattgcggg 840
cctccagctc atctttcacc tcacccctc cctcaagctt ggctttaatt atgctaagt 900
tgaggagaaa tgaataaata caagtga 928

```

<210> 33

<211> 762

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 4635039CB1

<400> 33

```

ccacatctgt cctctagaga atcccctgag agctccgttc ctcaacctgg actggacctg 60
gaggatcctc ttcttggtgg cagcagtcac aggagtccac tcccaagtgc agctggtgca 120
gtctggggct gaggtgagga agcctggggc ctcaagtgaag gtctcctgta aggttcttgg 180
ctacaccttc agcgaccact atattcactg ggttcgacag gccctggac aaggacttga 240
gtggatggga tggatcaacc ctaatagtgg tggcgcaagg tatgcacagg gctttcaggg 300
cttggtcacc atgaccaggg acacgtccat cagtacagcc tacttggagc tgcgcgggct 360

```

```

gagatctgac ggctcggccg tgtacttctg tgcgagacaa accacctcgt ctctgttagg 420
agatgctttt gatattctgg gccaagggac aatggtcacc gtctcttcag catccccgac 480
cagccccaag gtcttcccg cagacccag ccagatggga acgtgggcat 540
cgctgcttg gtccagggct tcttccccca ggagccactc agtgtgacct ggagcgaaac 600
ggaccagggc gtgaccgcca aaaaacttcc caccagcca ggatgccttc gggggactgt 660
aaaccacgag cagccagctg accctgcccgg gcaaaacagt gcctaaccgg gaaaattcgg 720
tgaaattgcc acgtgaagca ttaaaggatt cccagccagg gt 762

```

<210> 34

<211> 925

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 3240710CB1

<400> 34

```

cttctcacca tgaggctccc tgetcagctc ctggggctgc ttatgctctg gatacctgga 60
tcgagtgcgg acattgtgtt gaccagact ccactctctc tgtccgtcac cccgggacag 120
ccggcctcca tctcctgcaa gtottctgag agcctcctgc atactgatgg aaagacctat 180
ttgcattggg tcgtgcagaa ggcaggccag ccgccacaag tcctgatgta tgaagtctcc 240
aacgggttct ctggagtgcc agataggttc agtggcagcg ggtcaggcac agatttcaca 300
ctgaaaatca gccgggtgga ggctgaggat gttcggattt attactgcat gagaactata 360
caggttcctc caacgtggac gttcggccaa gggaccaagg tggaaatcaa acgaactgtg 420
gctgcaccat ctgtcttcat ctccccgcca tctgatgagc agttgaaatc tggaaactgcc 480
tctgttgtgt gcctgtgtaa taacttctat cccagagagg ccaaagtaca gtggaagggtg 540
gataacgccc tccaatcggg taactccag gagagtgtca cagagcagga cagcaaggac 600
agcacctaca gcctcagcag caccctgacg ctgagcaaag cagactacga gaaacacaaa 660
gtctacgcct gcgaagtcac ccatcagggc ctgagctcgc ccgtcacaaa gagcttcaac 720
aggggagagt gttagaggga gaagtgcgcc cactgtctcc tcagttccag cctgaccccc 780
tcccattcct tggcctctga ccccttttcc acaggggacc taccctatt gcggtcctcc 840
agctcatctt tcacctcacc cccctcctcc tccttggett taattatgct aatgttggag 900
gagaatgaat aaataaagtg aatcc 925

```

<210> 35

<211> 1584

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 4945813CB1

<400> 35

```

catggacctc ctgtgcaaga acatgaagca cctgtgggtc ttcctcctgc tgggtggcggc 60
tcccagatgg gtctgtccc agctgcaact gcaggagtcg ggcccaggat tgggtgaagcc 120
ttcggagacc ctgtccctca cctgcactgt ctctgggtggc tccatcagca gttataatca 180
ctactggggc tgggtccgcc agccccagg gaaggggctg gaatggattg ggagtatctt 240
ttatactggg aattcttact acaaccgctc cctcaagagt cgactcgcca tatcagtaga 300
cacgtccaag agccagctct ccctgaagct gagctctgtg accgcccgcg acacggccgt 360
gtattactgt gcgacagttc ctaaaaccgg gtctcgacca cgtggatata cctatgggtc 420
ctttgacttc tggggccagg gaaccctggt caccgtctcc tcagcatccc cgaccagccc 480

```

```

caaggtcttc ccgctgagcc tctgcagcac ccagccagat gggaacgtgg tcatcgctg 540
cctggtccag ggcttcttcc cccaggagcc actcagtgtg acctggagcg aaagcggaca 600
gggcgtgacc gccagaaact tcccacccag ccaggatgcc tccggggacc tgtacaccac 660
gagcagccag ctgaccctgc cggccacaca gtgcctagcc ggcaagtccg tgacatgcca 720
cgtgaagcac tacacgaatc ccagccagga tgtgactgtg ccctgcccag ttccttcaac 780
tccacctacc ccatctccct caactccacc taccatctt ccctcatgct gccacccccg 840
actgtcactg caccgaccgg ccctcgagga cctgctctta ggttcagaag cgaacctcac 900
gtgcacactg accggcctga gagatgcctc aggtgtcacc ttacacctga cgccctcaag 960
tggaagagc gctgttcaag gaccacctga gcgtgacctc tgtggtgct acagcgtgtc 1020
cagtgtcctg cggggtgtg ccgagccatg gaacctggg aagaccttca cttgactgc 1080
tgctacccc gagtccaaga ccccgctaac cgccacctc tcaaaatccg gaaacacatt 1140
cgggcccag gtccacctgc tgccgcccgc gtcggaggag ctggccctga acgagctggg 1200
gacgctgacg tgcctggcac gcggcttcag cccaaggac gtgctggttc gctgggtgca 1260
ggggtcacag gagctgcccc gcgagaagta cctgacttgg gcatcccggc aggagcccag 1320
ccagggcacc accaccttcg ctgtgaccag catactgcgc gtggcagccg aggactggaa 1380
gaagggggac accttctcct gcatggtggg ccacgagggc ctgccgctgg ccttcacaca 1440
gaagaccatc gaccgcttgg cgggtaaacc caccatgtc aatgtgtctg ttgtcatggc 1500
ggaggtggac ggcacctgct actgagccgc ccgctgtcc ccaccctga ataaactcca 1560
tgctcccca aaaaaaaaaa aaaa 1584

```

<210> 36
 <211> 804
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID NO: 4948957CB1

```

<400> 36
ctgagctaca acaggcaggc aggggcagca agatggtggt gcagaccag gtcttcattt 60
ctctgttgct ctggatctct gtccctgactg cagggtgccta cggggacatc gtgatgacct 120
agtctccaga ctccctggct gtgtctctgg gcgagagggc caccatcacc tgcaagtcca 180
gccagagtgt tttctacaac tccaacaata agaactactt agtttggtac cagcaaaagc 240
caggacagcc tcctaaaatg ctcatcttact gggcatctac ccgggaatcc ggggtccctg 300
accgattcag tggcagcggg tctgggacag atttcaactc caccatcagc agcctgcagg 360
ctgaagatgt ggcactttat tactgtcagc aatattttac tactccgtac acttttggcc 420
aggggaccag gctggagatc aaacgaactg tggctgcacc atctgtcttc atcttcccgc 480
catctgatga gcagttgaaa tctggaactg cctctgttgt gtgcctgctg aataacttct 540
atcccagaga ggccaaagta cagtggaagg tggataacgc cctccaatcg ggtaactccc 600
aggagagtgt cacagagcag gacagcaagg acagcaccta cagcctcagc agcacctga 660
cgctgagcaa agcagactac gagaaacaca aagtctacgc ctgcgaagtc acccatcagg 720
gcctgagctc gcccgtcaca aagagcttca acaggggaga gtgttagagg gagaagtgcc 780
cccacctgct cctcagttcc aggg 804

```

<210> 37
 <211> 878
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID NO: 4949649CB1

<400> 37

```

gcctgagagc acttagcggt catgagtggt cccaccatgg cctggatgat gcttctctctc 60
ggactccttg cttatggatc aggagtggat tctcagactg tggtgacca agagccatcg 120
ttatcagtgt cccctggagg gacagtcaca ctactttgtg gcttggcctc tgactcagtc 180
tctactaatt tcttccccac ctggtaccag cagaccccag gccaggctcc acgcacgctc 240
atctacagca caagcactcg ctcttctggg gtccctgatc gtttctctgg ctccatcctt 300
gggaacaaag ctgccctcac cattacgggg gccaggcag atgatgaatc tgattattac 360
tgtgcgcttt atatgggcag tggcatttct gtattcggcg gagggacca ggtgaccgtc 420
ctaggtcagc ccaaggctgc cccctcggtc actctgttcc cgccctcctc tgaggagctt 480
caagccaaca aggccacact ggtgtgtctc ataagtgact tctacccggg agccgtgaca 540
gtggcctgga aggcagatag cagccccgtc aaggcgagg tggagaccac cacacctcc 600
aaacaaagca acaacaagta cgcggccagc agctatctga gcctgacgcc tgagcagtgg 660
aagtcaccaca gaagtacag ctgccaggtc acgcatgaag ggagcaccgt ggagaagaca 720
gtggccctta cagaatgttc ataggttctc aacctcacc cccaccacg ggagactaga 780
gctgcaggat cccaggggag ggggtctctc tcccaccca aggcatacaag ccttctctcc 840
tgactcaat aaacctcaa taaatattct cattgtca 878

```

<210> 38

<211> 1680

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID NO: 5500302CB1

<400> 38

```

ctgggcgctg cggcggcagg aggaggacgg ggaaggacgg agccgagccg cggctgcctc 60
cctcgctcac tccctcggc actcgccgc cccctccctc cctccctcc cttccccggc 120
cccggtctct gccccggccc attcgctgtt ggtcttctg ctaggaggga tgtcgggttc 180
gtcgctgccc agcgccctgg cctctctcgt gttgctggtc tctggctccc tccctccagg 240
gccaggcgcc gctcagaacg ctgggtttgt caagtcgccc atgtcagaaa ctaagctcac 300
gggggacgcc tttgagctgt actgtgacgt ggtcgggagc cccacgccag agatccagt 360
gtggtacgca gaagtcaacc gggcagagtc tttcagacag ctgtgggacg gtgctcggaa 420
cgcgcgtgtc accgtaaaca cgcctacgg gtcaaacggc gtgagtgtgc tgagaataac 480
ccggctcacc ttggaggact ctgggactta cgagtgcagg gccagcaacg accccaagag 540
gaatgacttg aggcaaaacc cctccataac atggattcga gccaggcca ccataagcgt 600
ccttcagaag ccaaggattg tcaccagtga agaggtcatt attcgagaca gccctgttct 660
ccctgtcacc ctgcagtgtg acctcacctc cagctctcac acccttacat acagctactg 720
gacaaagaat ggggtggaac tgagtgccac tcgtaagaat gccagcaaca tggagtacag 780
gatcaataag ccgagagctg aggattcagg cgaataccac tgcgtatatc actttgtcag 840
cgctcctaaa gcaaacgcca ccattgaagt gaaagccgct cctgacatca ctggccataa 900
acggagttag aacaagaatg aagggcagga tgccactatg tattgcaagt cagttggcta 960
ccccaccca gactggatat ggcgcaagaa ggagaacggg atgcccattg acattgtcaa 1020
tacctctggc cgcttcttca tcatcaacaa ggaaaattac actgagttga acattgtgaa 1080
ctgcagatc acggaagacc ctggcgagta tgaatgtaat gccaccaacg ccattggctc 1140
cgctctgtt gtactgtcc tcagggtgcg gagccacctg gcccactct ggcttctctt 1200
gggaattctg gctgaaatta tcatccttgt ggtgatcatt gttgtgtatg agaagaggaa 1260
gaggccagat gaggttctct acgatgatga accagctgga ccaatgaaaa ccaactctac 1320
caacaatcac aaagataaaa acttgcgcca gagaaacaca aattaagtac tgcttacaat 1380
atcttttagt tcctgaaact ggtggcaaca tgacctgcta aaattttctg cttggacctc 1440
tttggttctc tcccctttca agtgagcaac accacaatga ctgtctaaag catgccttat 1500
ttagcctctc ctgtaagggt gatctagcca ggtacatttt aaacaatgct tcagtgtaga 1560
aggtgtaaac tattttgggc ttgatgtgct gtgaatgttg cttttttttt ccttggtaaa 1620

```


atattttaa at tagaagtgaa aaggtcctct gaggatcaga ttcatgcatg cgccattttt 1680

DECLARATION AND POWER OF ATTORNEY FOR UNITED STATES PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,
and

I believe that I am the original, first and sole inventor (if only one name is listed below)
or an original, first and joint inventor (if more than one name is listed below) of the subject
matter which is claimed and for which a United States patent is sought on the invention entitled

IMMUNOGLOBULIN SUPERFAMILY PROTEINS

the specification of which:

 / is attached hereto.

 / was filed on as application Serial No. and if this box
contains an X /, was amended on .

 X / was filed as Patent Cooperation Treaty international application No. PCT/US99/27566
on November 19, 1999, if this box contains an X /, was amended on under Patent Cooperation
Treaty Article 19 on 2001, and if this box contains an X /, was amended on .

I hereby state that I have reviewed and understand the contents of the above-identified
specification, including the claims, as amended by any amendment referred to above.

I acknowledge my duty to disclose information which is material to the examination of
this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim the benefit under Title 35, United States Code, §119 or §365(a)-(b) of any
foreign application(s) for patent or inventor's certificate indicated below and of any Patent
Cooperation Treaty international applications(s) designating at least one country other than the
United States indicated below and have also identified below any foreign application(s) for
patent or inventor's certificate and Patent Cooperation Treaty international application(s)
designating at least one country other than the United States for the same subject matter and
having a filing date before that of the application for said subject matter the priority of which is
claimed:

| Country | Number | Filing Date | Priority Claimed |
|-----------------------------|-----------------------------|-----------------------------|--|
| <u> </u> | <u> </u> | <u> </u> | <u> </u> / <u> </u> Yes <u> </u> / <u> </u> No |
| <u> </u> | <u> </u> | <u> </u> | <u> </u> / <u> </u> Yes <u> </u> / <u> </u> No |

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

| Application Serial No. | Filed | Status (Pending, Abandoned, Patented) |
|---------------------------|-------------------|--|
| 60/183,028 | November 19, 1998 | Expired |
| 60/113,635 | December 22, 1998 | Expired |
| 60/128,194 | April 7, 1999 | Expired |

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in said prior application(s) in the manner required by the first paragraph of Title 35, United States Code §112, I acknowledge my duty to disclose material information as defined in Title 37 Code of Federal Regulations, §1.56(a) which occurred between the filing date(s) of the prior application(s) and the national or Patent Cooperation Treaty international filing date of this application:

| Application Serial No. | Filed | Status (Pending, Abandoned, Patented) |
|---------------------------|-------|--|
|---------------------------|-------|--|

I hereby appoint the following:

| | |
|----------------------|------------------------|
| Lucy J. Billings | Reg. No. <u>36,749</u> |
| Michael C. Cerrone | Reg. No. <u>39,132</u> |
| Diana Hamlet-Cox | Reg. No. <u>33,302</u> |
| Richard C. Ekstrom | Reg. No. <u>37,027</u> |
| Barrie D. Greene | Reg. No. <u>46,740</u> |
| Matthew R. Kaser | Reg. No. <u>44,817</u> |
| Lynn E. Murry | Reg. No. <u>42,918</u> |
| Shirley A. Recipon | Reg. No. <u>47,016</u> |
| Susan K. Sather | Reg. No. <u>44,316</u> |
| Michelle M. Stempien | Reg. No. <u>41,327</u> |
| David G. Streeter | Reg. No. <u>43,168</u> |
| Stephen Todd | Reg. No. <u>47,139</u> |
| Christopher Turner | Reg. No. <u>45,167</u> |
| P. Ben Wang | Reg. No. <u>41,420</u> |


respectively and individually, as my patent attorneys and/or agents, with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith. Please address all communications to:

**LEGAL DEPARTMENT
 INCYTE GENOMICS, INC.
 3160 PORTER DRIVE, PALO ALTO, CA 94304**


TEL: 650-855-0555 FAX: 650-849-8886 or 650-845-4166

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.


First Joint Inventor:

Full name: Henry Yue
Signature: 
Date: March 02, 2001
Citizenship United States of America
Residence: Sunnyvale, California CA.
P.O. Address: 826 Lois Avenue
Sunnyvale, California 94087


Second Joint Inventor:

Full name: Y. Tom Tang
Signature: 
Date: February 27, 2001
Citizenship ~~People's Republic of China~~ USA
Residence: San Jose, California CA.
P.O. Address: 4230 Ranwick Court
San Jose, California 95118


3-00
Third Joint Inventor:

Full name: Neil C. Corley
 Signature: 
 Date: MAR 5, 2001
 Citizenship: United States of America
 Residence: Castro Valley, California CA.
 P.O. Address: 20426 Crow Creek Road
Castro Valley, California 94552

4-00
Fourth Joint Inventor:


Full name: Karl J. Guegler
 Signature: 
 Date: 02/02, 2001
 Citizenship: Switzerland
 Residence: Menlo Park, California CA.
 P.O. Address: 1048 Oakland Avenue
Menlo Park, California 94025

5-00
Fifth Joint Inventor:


Full name: Gina A. Gorgone
 Signature: 
 Date: Feb 5, 2001
 Citizenship: United States of America
 Residence: Boulder Creek, California CA.
 P.O. Address: 1253 Pinecrest Drive
Boulder Creek, California 95006

FOOTNOTES - SEE PAGES 4-5


6-00
Sixth Joint Inventor:

Full name: Mariah R. Baughn
 Signature: 
 Date: February 12, 2001
 Citizenship: United States of America
 Residence: San Leandro, California CA.
 P.O. Address: 14244 Santiago Road
San Leandro, California 94577

7-00
Seventh Joint Inventor:


Full name: Dyung Aina M. Lu
 Signature: 
 Date: March 22, 2001
 Citizenship: United States of America
 Residence: San Jose, California CA.
 P.O. Address: 233 Coy Drive
San Jose, California 95123

8-00
Eighth Joint Inventor:


Full name: Preeti Lal
 Signature: 
 Date: FEBRUARY, 16, 2001
 Citizenship: India
 Residence: Santa Clara, California CA.
 P.O. Address: P.O. Box 5142
Santa Clara, California 95056

FOOTNOTES - SEE PAGES 5-8

Ninth Joint Inventor:

Full name: Jennifer L. Hillman
Signature: 
Date: February 16, 2001
Citizenship: United States of America
Residence: Mountain View, California CA
P.O. Address: 230 Monroe Drive, #17
Mountain View, California 94040

Tenth Joint Inventor:

Full name: Junming Yang
Signature: 
Date: _____, 2001
Citizenship: China
Residence: San Jose, CA
Castro Valley, California CA
P.O. Address: 7125 Bark Lane
San Jose, California 95129